

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
19 December 2002 (19.12.2002)

PCT

(10) International Publication Number  
WO 02/101059 A2

- (51) International Patent Classification<sup>7</sup>: C12N 15/82, C07K 14/415, A01H 5/00
- (21) International Application Number: PCT/EP02/06265
- (22) International Filing Date: 7 June 2002 (07.06.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
101 28 363.6 12 June 2001 (12.06.2001) DE
- (71) Applicant: BAYER CROPSCIENCE GMBH [DE/DE];  
Brüningstrasse 50, 65929 Frankfurt (DE).
- (72) Inventors: UWER, Ursula; Hasenheide 49, 10967 Berlin (DE). FROHBERG, Claus; Apoldaer Strasse 14, 12249 Berlin (DE). PILLING, Jens; Melchiorstrasse 36, 50670 Köln (DE). LANDSCHÜTZE, Volker; Gardeschützenweg 51, 12203 Berlin (DE).
- (74) Agent: VOSSIUS & PARTNER; Siebertstrasse 4, 81675 Munich (DE).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: TRANSGENIC PLANTS SYNTHESISING HIGH AMYLOSE STARCH

(57) Abstract: Transgenic plant cells and plants are described which synthesise a starch which is modified in comparison with corresponding wild type plant cells and plants. The plant cells and plants described show a reduced activity of R1, BEI and BEII proteins. Furthermore, modified starches as well as methods for their production are described.

WO 02/101059 A2

### **Transgenic plants synthesising high amylose starch**

The present invention relates to genetically modified plant cells and plants wherein the genetic modification leads to the reduction of the activity of R1 and BEI and BEII proteins in comparison with corresponding plant cells of wild type plants that have not been genetically modified. Furthermore, the present invention relates to means and methods for the production thereof. Plant cells and plants of that type synthesise a modified starch characterised in that it has an amylose content of at least 75% and - in comparison with starch of corresponding wild type plants which have not been genetically modified - a reduced phosphate content and/or a modified distribution of the side chains and/or an increased gel strength in the texture analyser and/or a modified starch granule morphology and/or a modified average starch granule size. Thus, the present invention also relates to starch that can be synthesised by the plant cells and plants of the invention as well as methods for the production of this starch.

With regard to the increasing importance of plant ingredients as renewable raw material sources in the past few years, one of the problems of research in the field of biotechnology is to endeavour adjustment of these raw materials to the requirements of the processing industry. For allowing an application of renewable raw materials in as many as fields as possible, it is furthermore necessary to achieve a great variety of substances.

Apart from oils, fats and proteins, polysaccharides represent the essential renewable raw materials from plants. Among the polysaccharides, starch plays a central role beside cellulose. It is one of the most important storage substances in higher plants. For allowing as wide an application of starch as possible, it seems desirable to provide plants which are able to synthesise modified starch that is particularly suitable for different purposes. One possibility of providing such plants is - apart from cultivating - the purposeful genetic modification of the starch metabolism of starch-producing plants by genetic engineering.

The polysaccharide starch is a polymer of chemically uniform basic building blocks - the glucose molecules. It is, however, a very complex mixture of different molecule forms which differ with regard to their polymerisation degree and the occurrence of branchings of the glucose chains. Thus, starch is not a uniform raw material. There are two chemically different components of starch: the amylose and the amylopectin. In plants typically used for the production of starch, such as e.g. maize, wheat or potato, the synthesised starch consists of about 20% - 30% of amylose starch and of about 70% - 80% of amylopectin starch.

Amylose was considered a linear polymer for a long time, consisting of  $\alpha$ -1,4-glycosidically bound  $\alpha$ -D-glucose monomers. In recent studies, however, the presence of about 0.1%  $\alpha$ -1,6-glycosidic branching points has been proven (Hizukuri and Takagi, Carbohydr. Res. 134, (1984), 1-10; Takeda et al., Carbohydr. Res. 132, (1984), 83-92).

As a rule, the complete separation of the amylose from the amylopectin is very difficult so that the quality of the amylose strongly depends on the type of the separation method chosen.

There are different methods for the determination of the amylose content. Some of these methods are based on the iodine-binding capacity of the amylose which can be determined potentiometrically (Banks & Greenwood, in W. Banks & C.T. Greenwood, Starch and its components (page 51-66), Edinburgh, Edinburgh University Press), amperometrically (Larson et al., Analytical Chemistry 25(5), (1953), 802-804) or spectrophotometrically (Morrison & Laignelet, J. Cereal Sc. 1, (1983), 9-20). The determination of the amylose content can also be carried out calorimetrically by means of DSC (differential scanning calorimetry) measurements (Kugimiya & Donovan, Journal of Food Science 46, (1981), 765-770; Sievert & Holm, Starch/Stärke 45 (4), (1993), 136-139). Moreover, it is possible to determine the amylose content by using the SEC (size exclusion chromatography) of native or debranched starch. This method was particularly recommended for the determination of the amylose content of genetically modified starches (Gérard et al., Carbohydrate Polymers 44, (2001), 19-27).

The choice of the analysis method used for the determination of the amylose content of a starch has a crucial influence on the size of the amylose figures determined as

could be shown by various studies (Shi et al., J. Cereal Science 27, (1998), 289-299; Gérard et al., Carbohydrate Polymers 44, (2001), 19-27).

In contrast to the amylose, the amylopectin is branched to a larger degree and exhibits about 4% branching points which occur due to the presence of additional  $\alpha$ -1,6-glycosidic linkings. The amylopectin is a complex mixture of glucose chains branched differently.

A further essential difference between amylose and amylopectin is the molecular weight. While amylose - depending on the origin of the starch - has a molecular weight of  $5 \times 10^5$  -  $10^6$  Da, the molecular weight of amylopectin is between  $10^7$  and  $10^8$  Da. Both macromolecules can be differentiated from each other by their molecular weight and their different physico-chemical properties, which can be made apparent in the simplest way by their different iodine-binding properties.

The functional properties of the starch are strongly influenced - apart from the amylose/amylopectin ratio and the phosphate content - by the molecular weight, the pattern of the side chain distribution, the content of ions, the lipid and protein content, the average starch granule size and the starch granule morphology etc. Important functional properties to be mentioned are, for example, the solubility, the retrogradation behaviour, the water binding capacity, the film formation properties, the viscosity, the pasting properties, the freeze-thaw-stability, the acid stability, the gel strength etc. The starch granule size, too, can be important for different applications.

The ratio of amylopectin and amylose has a strong influence on the physico-chemical properties of the starches and, thus, on the possible applications of these starches. Since methods for the separation of these two components are very time-consuming and costly, such methods are no longer used on a large technical scale (Yound, A.H. in: Starch Chemistry and Technology. Eds. R. L. Whistler, J. N. BeMiller and E. F. Paschall. Academic Press, New York, 1984, 249-283). For a plurality of applications it would be desirable to have starches at disposal which still contain only one of the two polymers or at least one of the two starch components in an enriched form.

So far, both mutants and plants produced by genetic engineering have been described which, in comparison with corresponding wild type plants, exhibit a modified amylopectin/amylose ratio.

For example, the so-called "waxy" mutant from maize exhibiting a mutation in the gene encoding the starch granule bound starch synthase I (abbreviated: GBSSI) (Akasaka and Nelson, J. Biol. Chem., 241, (1966), 2280-2285; Shure et al., Cell 35 (1983), 225-233), produces a starch essentially consisting of amylopectin. For potato, genotypes were produced both by means of chemical mutagenesis of a haploid line (Hovenkamp-Hermelink et al., Theor. Appl. Genet., 225, (1987), 217-221) and by means of antisense inhibition of the GBSSI-gene, whose starches essentially consist of amylopectin starch. In comparison with starches of the corresponding wild type plants, such waxy potato starches do not exhibit any differences with regard to phosphate content, the morphology of the starch granule or the ion content (Visser et al., Starch/Stärke, 49, (1997), 438-443).

Furthermore, maize mutants are commercially available which exhibit starches with amylose contents of about 50% or about 70% (amylose content determined by potentiometric determination of the iodine-binding capacity) and which are designated Hylon V® or HylonVII® (National Starch and Chemical Company, Bridgewater, NJ, USA). Moreover, also maize hybrids have been described which synthesise so-called "low amylopectin starch" (LAPS) and exhibit a content of high molecular ("high mol weight") amylopectin of about 2.5% and an amylose content of about 90% (potentiometric determination of the iodine-binding capacity) (Shi et al., J. Cereal Science 27, (1998), 289-299).

Furthermore, transgenic potato plants have been described which, due to the antisense-inhibition of the branching enzyme I (= BEI) and the branching enzyme II (= BEII) gene, synthesise a potato starch which exhibits an amylose content of up to 75% by colorimetric determination of the amylose content according to the method described by Morrison and Laignelet (J. Cereal Sci. 1, (1983), 9-20) (Schwall et al., Nature Biotechn. 18, (2000), 551-554). These potato starches are characterised by a phosphate content of the starch which is up to 6 times higher compared to corresponding wild type plants. Furthermore, the international patent application WO 97/11188 describes transgenic potato plants which, due to their antisense inhibition

of the R1 gene and the BEI gene synthesise a starch with an amylose content of more than 70%, the amylose content having been determined according to the method by Hovenkamp & Hermelink (Potato Research 31, (1988), 241-246).

Transgenic potato plant cells and potato plants synthesising a starch having an amylose content of more than 75% (colorimetric determination of the amylose content according to Hovenkamp & Hermelink (Potato Research 31, (1988), 241-246) and, at the same time, a reduced phosphate content in comparison with corresponding wild type plants have not been described in the state of the art so far. The same applies to the potato starches that can be isolated from these potato plant cells and plants and to methods for the production of such starches. However, the provision of such starches is desirable since their physico-chemical properties can be expected to be advantageously useful for various industrial applications.

Thus, the technical problem underlying the present invention is to provide plant cells and plants synthesising starch which has an amylose content of more than 75% (colorimetric determination of the amylose content according to Hovenkamp & Hermelink (Potato Research 31, (1988), 241-246) and a reduced phosphate content in comparison with the phosphate content of starch from corresponding wild type plant cells and plants that have not been genetically modified, as well as to provide such starch which differs from the starches described in the state of the art in its structural and/or functional properties and is, thus, more suitable for general and/or specific purposes.

This technical problem has been solved by providing the embodiments characterised in the claims.

Thus, the present invention relates to a transgenic plant cell which is genetically modified, wherein the genetic modification leads to a reduction of the activity of one or more R1 proteins occurring endogenously in the plant cell and to the reduction of the activity of one or more BEI proteins occurring endogenously in the plant cell and to the reduction of the activity of one or more BEII proteins occurring endogenously

in the plant cell in comparison with corresponding plant cells of wild type plants, the cells not being genetically modified.

The genetic modification may be any genetic modification leading to a reduction of the activity of one or more R1 proteins occurring endogenously in the plant cell and to the reduction of the activity of one or more BEI proteins occurring endogenously in the plant cell and to the reduction of the activity of one or more BEII proteins occurring endogenously in the plant cell in comparison with corresponding plant cells of wild type plants, the cells not being genetically modified.

In this context, the term "transgenic" means that the plant cells of the invention differ in their genetic information from corresponding plant cells which are not genetically modified due to a genetic modification, in particular the introduction of one or more foreign nucleic acid molecules.

In this context, the term "genetically modified" means that the plant cell is modified in its genetic information due to the introduction of one or more foreign nucleic acid molecules and that the presence and/or the expression of the foreign nucleic acid molecule/s lead/s to a phenotypic modification. In this context, phenotypic modification preferably relates to a measurable modification of one or more functions of the cells. Genetically modified plants cells of the invention, for example, exhibit a reduction of the expression of one or more R1 genes occurring endogenously in the plant cell and a reduction of the expression of one or more BEI genes occurring endogenously in the plant cell and a reduction of the expression of one or more BEII genes occurring endogenously in the plant cell in comparison with corresponding plant cells of wild type plants, the cells not being genetically modified, and/or a reduction of the activity of one or more R1 proteins occurring endogenously in the plant cell and a reduction of the activity of one or more BEI proteins occurring endogenously in the plant cell and a reduction of the activity of one or more BEII proteins occurring endogenously in the plant cell in comparison with corresponding plant cells of wild type plants, the cells not being genetically modified.

Within the meaning of the present invention, the term "reduction of the activity" means a reduction of the expression of endogenous genes encoding R1, BEI and/or BEII proteins and/or a reduction of the amount of R1, BEI and/or BEII proteins in the cells and/or a reduction of the enzymatic activity of the R1, BEI and/or BEII proteins in the cells.

In the context of the present invention, the term "reduction of expression" refers to a reduction of the amount of transcripts of the respective endogenous gene in a plant cell of the invention as compared to a corresponding wild-type plant cell. The reduction of the expression can, for instance, be determined by measuring the amount of transcripts encoding R1, BEI or BEII proteins, e.g. by means of Northern blot analysis or RT-PCR. In this context, a reduction preferably means a reduction of the amount of transcripts in comparison with corresponding cells that have not been genetically modified by at least 50%, in particular by at least 70%, more preferably by at least 85% and most preferably by at least 95%.

The reduction of the amount of R1, BEI and/or BEII proteins can, for instance, be determined by means of Western blot analysis. In this context, a reduction preferably means a reduction of the amount of R1, BEI and/or BEII proteins in comparison with corresponding cells which have not been genetically modified by at least 50%, in particular by at least 70%, more preferably by at least 85% and most preferably by at least 95%.

Methods for determining the reduction of the enzymatic activity of the R1, BEI and BEII proteins are known to the person skilled in the art and will be described further below for each protein individually. In the context of the present invention, the term "R1 protein" relates to proteins which have been described, for example, in Lorberth et al. (Nature Biotech. 16, (1998), 473-477) and in the international patent applications WO 98/27212, WO 00/77229, WO 00/28052 and which have the following characteristics. Important characteristics of R1 proteins are i) their amino acid sequence (see, for example, GenBank Acc. No. A61831, Y09533); ii) their localisation in the plastides of plant cells; iii) their ability to influence the degree of



phosphorylation of starch in plants.

Further, the term "R1 protein" refers to a protein catalysing the phosphorylation of starch in dikinase-type reaction in which three substrates, an  $\alpha$ -polyglucan, ATP and  $H_2O$  are converted into three products, an  $\alpha$ -polyglucan-P, AMP and orthophosphate (Ritte et al., PNAS 99(10) (2002), 7166-7171).

The inhibition of the R1 gene encoding an R1 protein from potato in transgenic potato plants, for example, leads to a reduction of the phosphate content of the starch which can be isolated from the potato tuber. Moreover, Lorberth et al. were able to demonstrate that the R1 protein from *Solanum tuberosum* is able to phosphorylate bacterial glycogen when the corresponding R1 cDNA is expressed in *E. coli* (Lorberth et al., Nature Biotech. 16, (1998), 473-477).

Ritte et al. (Plant J. 21, (2000), 387-391) were able to show that the R1 protein from *Solanum tuberosum* in potato plants binds to starch granules in a reversible way, wherein the strength of the binding to the starch granule depends on the metabolic status of the plant. In potato plants, the protein is mainly present in starch granule bound form in leaves that have been kept in the dark. After exposing the leaves to light, however, the protein is mainly present in the soluble form which is not bound to the starch granule.

Furthermore, the inhibition of the expression of the R1 gene from potato in transgenic potato plants or in the tubers thereof leads to a reduction of the so-called "cold-induced-sweetenings" (Lorberth et al., Nature Biotech. 16, (1998), 473-477).

In the context of the present invention, the term "R1 protein" also relates to proteins which exhibit a significant homology (identity) of at least 60%, preferably of at least 80%, more preferably of at least 90% to the amino acid sequence stated under SEQ ID NO: 6 or under the GenBank Acc. No. Y09533 or A61831, and which are able to modify the degree of phosphorylation of polysaccharides such as, for example, starch and/or glycogen. Preferably, the R1 protein originates from potato (GenBank Acc. No. Y09533 or A61831).

Preferably, an R1 protein, as addressed in the embodiments of the present invention, is encoded by a nucleic acid molecule which hybridises, advantageously

under stringent conditions, with the nucleic acid molecule having the nucleotide sequences shown under SEQ ID NO: 5 and which encodes a polypeptide having the activity of an R1 protein.

Within the present invention the term "hybridization" means hybridization under conventional hybridization conditions (also referred to as "low stringency conditions"), preferably under stringent conditions (also referred to as "high stringency conditions"), as for instance described in Sambrook and Russell (2001), Molecular Cloning, A Laboratory Manual, CSH Press, Cold Spring Harbour, NY, USA. Within an especially preferred meaning the term "hybridization" means that hybridization occurs under the following conditions:

Hybridization buffer:	2 x SSC; 10 x Denhardt solution (Fikoll 400 + PEG + BSA; ratio 1:1:1); 0.1% SDS; 5 mM EDTA; 50 mM Na <sub>2</sub> HPO <sub>4</sub> ; 250 µg/ml of herring sperm DNA; 50 µg/ml of tRNA; or 0.25 M of sodium phosphate buffer, pH 7.2; 1 mM EDTA 7% SDS
Hybridization temperature T	= 60°C
Washing buffer:	2 x SSC; 0.1% SDS
Washing temperature T	= 60°C.

Nucleic acid molecules which hybridize with a nucleic acid molecule having the nucleotide sequence shown under SEQ ID NO: 5 can, in principle, encode a R1 protein from any organism expressing such a protein.

Such hybridizing nucleic acid molecules can for instance be isolated from genomic libraries or cDNA libraries of plants. Alternatively, they can be prepared by genetic engineering or chemical synthesis.

Such nucleic acid molecules may be identified and isolated with the use of a nucleic acid molecule encoding an R1 protein as disclosed herein or parts of such a molecule or reverse complements of such a molecule, for instance by hybridization according to standard methods (see for instance Sambrook and Russell (2001), Molecular Cloning. A Laboratory Manual, CSH Press, Cold Spring Harbor, NY,

USA).

Nucleic acid molecules possessing the same or substantially the same nucleotide sequence as indicated in SEQ ID NO: 5 or parts thereof can, for instance, be used as hybridization probes. The fragments used as hybridization probes can also be synthetic fragments which are prepared by usual synthesis techniques, and the sequence of which substantially coincides with that of a nucleic acid molecule specifically described herein.

The hybridizing nucleic acid molecules also comprise fragments, derivatives and allelic variants of the nucleic acid molecule having the nucleotide sequence shown under SEQ ID NO: 5. Herein, fragments are understood to mean parts of the nucleic acid molecules which are long enough to encode an R1 protein. In this connection, the term derivative means that the sequences of these nucleic acid molecules differ from the sequence of an above-described nucleic acid molecule in one or more positions and show a high degree of homology to such a sequence. In this context, homology means a sequence identity of at least 40%, in particular an identity of at least 60%, preferably of at least 65%, more preferably of at least 70%, even more preferably of at least 80%, in particular of at least 85%, furthermore preferred of at least 90% and particularly preferred of at least 95%. Most preferably homology means a sequence identity of at least  $n\%$ , wherein  $n$  is an integer between 40 and 100, i.e.  $40 \leq n \leq 100$ . Deviations from the above-described nucleic acid molecules may have been produced, e.g., by deletion, substitution, insertion and/or recombination.

Preferably, the degree of homology is determined by comparing the respective sequence with the nucleotide sequence of the coding region of SEQ ID No: 5. When the sequences which are compared do not have the same length, the degree of homology preferably refers to the percentage of nucleotide residues in the shorter sequence which are identical to nucleotide residues in the longer sequence. The degree of homology can be determined conventionally using known computer programs such as the ClustalW program (Thompson et al., Nucleic Acids Research 22 (1994), 4673-4680) distributed by Julie Thompson (Thompson@EMBL-

Heidelberg.DE) and Toby Gibson (Gibson@EMBL-Heidelberg.DE) at the European Molecular Biology Laboratory, Meyerhofstrasse 1, D 69117 Heidelberg, Germany. ClustalW can also be downloaded from several websites including IGBMC (Institut de Génétique et de Biologie Moléculaire et Cellulaire, B.P.163, 67404 Illkirch Cedex, France; <ftp://ftp-igbmc.u-strasbg.fr/pub/>) and EBI (<ftp://ftp.ebi.ac.uk/pub/software/>) and all sites with mirrors to the EBI (European Bioinformatics Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge CB10 1SD, UK).

When using ClustalW program version 1.8 to determine whether a particular sequence is, for instance, 90% identical to a reference sequence according to the present invention, the settings are set in the following way for DNA sequence alignments:

KTUPLE=2, TOPDIAGS=4, PAIRGAP=5, DNAMATRIX:IUB, GAOPEN=10, GAPEXT=5, MAXDIV=40, TRANSITIONS: unweighted.

For protein sequence alignments using ClustalW program version 1.8 the settings are the following: KTUPLE=1, TOPDIAG=5, WINDOW=5, PAIRGAP=3, GAOPEN=10, GAPEXTEND=0.05, GAPDIST=8, MAXDIV=40, MATRIX=GONNET, ENDGAPS(OFF), NOPGAP, NOHGAP.

Homology, moreover, means that there is a functional and/or structural equivalence between the corresponding nucleic acid molecules or proteins encoded thereby. Nucleic acid molecules which are homologous to one of the above-described molecules and represent derivatives of these molecules are generally variations of these molecules which represent modifications having the same biological function. They may be either naturally occurring variations, for instance sequences from other microorganisms, or mutations, and said mutations may have formed naturally or may have been produced by deliberate mutagenesis. Furthermore, the variations may be synthetically produced sequences. The allelic variants may, e.g., be naturally occurring variants or synthetically produced variants or variants produced by recombinant DNA techniques.

The proteins encoded by the different variants of the nucleic acid molecule having the nucleotide sequence shown under SEQ ID NO: 5 possess certain characteristics they have in common. These include for instance enzymatic activity, molecular weight, immunological reactivity, conformation, etc., and physical properties, such as

for instance the migration behavior in gel electrophoreses, chromatographic behavior, sedimentation coefficients, solubility, spectroscopic properties, stability, pH optimum, temperature optimum etc.

In the context of the present invention, the term "R1 gene" relates to a nucleic acid molecule (e.g. cDNA, DNA) encoding an "R1 protein" as described above. Nucleic acid molecules encoding R1 proteins have been described for various plants such as, e.g. maize (WO 98/27212 A1), rice (WO 00/28052 A1) and wheat (WO 00/77229 A1). Preferably, the R1 gene originates from potato, an R1 cDNA from potato with the nucleotide sequence stated under SEQ ID NO: 5 or GenBank Acc. No. Y09533 or A61831 is particularly preferred.

In the context of the present invention, the term "branching enzyme" or "BE protein" ( $\alpha$ -1,4-glucan:  $\alpha$ -1,4-glucan 6-glycosyltransferase, E.C. 2.4.1.18) relates to a protein catalysing a transglycosylation reaction, wherein  $\alpha$ -1,4-linkings of an  $\alpha$ -1,4-glucan donor are hydrolysed and the  $\alpha$ -1,4-glucan chains released thereby are transferred to an  $\alpha$ -1,4-glucan acceptor chain and are thereby converted into  $\alpha$ -1,6-linkings. Within the meaning of the present invention, the term "BE gene" is a gene encoding a "BE protein".

In the context of the present invention, the term "BEI protein" relates to a branching enzyme (= BE) of the isoform I, preferably the BEI protein originates from potato plants.

The designation of the isoforms follows the nomenclature suggested by Smith-White and Preiss (Smith-White and Preiss, *Plant Mol. Biol. Rep.* 12 (1994), 67-71; Larsson et al., *Plant Mol. Biol.* 37, (1998), 505-511). This nomenclature assumes that all branching enzymes exhibiting a higher homology (identity) on the amino acid level to the BEI protein from maize having the amino acid sequence shown under SEQ ID NO: 9 (GenBank Acc. No. D11081; Baba et al., *Biochem. Biophys. Res. Commun.* 181 (1), (1991), 87-94; Kim et al., *Gene* 216, (1998), 233-243) than to the BEII protein from maize having the amino acid sequence shown under SEQ ID NO: 10 (GenBank Acc. No. AF072725, U65948) are designated branching enzymes of the

isoform I or, in short, BEI proteins.

In the context of the present invention, the term "BEI gene" relates to a nucleic acid molecule (e.g. cDNA, DNA) encoding a "BEI protein", preferably a BEI protein from potato plants. Such nucleic acid molecules have been described for numerous plants, for example for maize (GenBank Acc. No. D11081, AF 072724), rice (GenBank Acc. No. D11082), pea (GenBank Acc. No. X80010) and potato. Different forms of the BEI gene or the BEI protein from potato have been described, for example, by Khoshnoodi et al. (Eur. J. Biochem. 242 (1) (1996), 148-155), GenBank Acc. No. Y08786 and by Kossmann et al. (Mol. Gen. Genet. 230 (1991), 39-44). In potato plants, the BEI gene is expressed mainly in the tubers and hardly in the leaves (Larsson et al., Plant Mol. Biol. 37, (1998), 505-511).

Preferably, a BEI protein, as addressed in the embodiments of the present invention, is encoded by a nucleic acid molecule which hybridises, advantageously under stringent conditions, with the nucleic acid molecule having the nucleotide sequence shown under SEQ ID NO: 7 and which encodes a polypeptide having branching enzyme activity.

The definition for the term "hybridisation" as defined above in connection with R1 proteins applies equally for the definition of nucleic acid molecules hybridising with the nucleic acid molecule having the nucleotide sequence of SEQ ID NO: 7. Preferably, a BEI protein as referred to herein displays a sequence identity of at least 60%, in particular of at least 75%, preferably of at least 85%, more preferably at least 90% and even more preferably at least 95% to the amino acid sequence depicted under SEQ ID NO: 8.

In the context of the present invention, the term "BEII protein" relates to a branching enzyme (= BE) of the isoform II, preferably it originates from potato plants. Within the meaning of the present invention, all enzymes exhibiting a higher homology (identity) on the amino acid level to the BEII protein from maize (GenBank Acc. No. AF072725, U65948) than to the BEI protein from maize (GenBank Acc. No. D 11081, AF 072724) are to be designated BEII protein.

In the context of the present invention, the term "BEII gene" relates to a nucleic acid molecule (e.g. cDNA, DNA) encoding a "BEII protein", preferably a BEII protein from potato plants. Such nucleic acid molecules have been described for numerous plants, for example, for potato (GenBank Acc. No. AJ000004, AJ011888, AJ011889, AJ011885, AJ011890), maize (AF072725, U65948), barley (AF064561), rice (D16201) and wheat (AF286319). In potato plants, the BEII gene is expressed mainly in the leaves and hardly in the tubers (Larsson et al., Plant Mol. Biol. 37, (1998), 505-511).

Preferably, a BEII protein, as addressed in the embodiments of the present invention, is encoded by a nucleic acid molecule which hybridises, advantageously under stringent conditions, with the nucleic acid molecule having the nucleotide sequence shown under SEQ ID NO: 9 and which encodes a polypeptide having branching enzyme activity.

The definition for the term "hybridisation" as defined above in connection with R1 proteins applies equally for the definition of nucleic acid molecules hybridising with the nucleic acid molecule having the nucleotide sequence of SEQ ID NO: 3. Preferably, a BEII protein as referred to herein displays a sequence identity of at least 60%, in particular of at least 75%, preferably of at least 85%, more preferably at least 90% and even more preferably at least 95% to the amino acid sequence depicted under SEQ ID NO: 4.

In a preferred embodiment of the present invention, the genetic modification of the transgenic plant cell of the invention is the introduction of one or more foreign nucleic acid molecules the presence and/or expression of which leads to a reduction of the activity of R1 and BEI and BEII proteins in comparison with corresponding plant cells of wild type plants, the cells not being genetically modified.

Preferably, this reduction of activity is achieved by inhibiting the expression of the endogenous genes encoding R1 proteins, BEI proteins and BEII proteins.

The production of the plant cells of the invention can be achieved by different methods known to the person skilled in the art, e.g. by methods leading to an inhibition of the expression of endogenous genes encoding an R1, BEI or BEII protein. These include, for example, the expression of a corresponding antisense RNA, the provision of molecules or vectors mediating a co-suppression effect, the expression of a ribozyme constructed accordingly which specifically cleaves transcripts encoding an R1, BEI or BEII protein or the so-called "in-vivo mutagenesis". Furthermore, the reduction of the R1 and/or the BEI and/or the BEII activity in the plant cells may also be caused by the simultaneous expression of sense and antisense RNA molecules of the target gene to be repressed, preferably of the R1 and/or the BEI and/or the BEII gene, a technique which is commonly referred to as RNA interference (RNAi) (Bosher and Labouesse, *Nature Cell Biology* 2, (2000), E31-E36; Waterhouse et al., *PNAS* 95, (1998), 13959-13964). Furthermore, by the use of double-stranded RNA molecules comprising promoter sequences, a transcriptional inactivation of the promoter can be achieved.

These and further methods for reducing the activity of proteins will be described in more detail below. All these methods are based on the introduction of one or more foreign nucleic acid molecules into the genome of plant cells.

Within the context of the present invention, the term "foreign nucleic acid molecule" is understood to be a molecule which either does not occur naturally in corresponding plant cells or which does not occur naturally in the plant cells in the concrete spatial order or which is located at a position in the genome of the plant cell at which it does not occur naturally. The foreign nucleic acid molecule preferably is a recombinant molecule which consists of various elements the combination or specific spatial order of which does not occur naturally in plant cells.

The foreign nucleic acid molecule can, for instance, be a so-called "triple construct" which is understood to be a single vector for plant transformation which contains both the genetic information for inhibiting the expression of one or more endogenous R1 genes and for inhibiting the expression of one or more BEI and BEII genes or the presence or expression of which leads to the reduction of the activity of one or more R1, BEI and BEII proteins.



In another embodiment, the foreign nucleic acid molecule may be a so-called "double construct" which is understood to be a vector for plant transformation which contains the genetic information for inhibiting the expression of two of the three target genes (R1, BEI, BEII gene) or the presence or expression of which leads to the reduction of the activity of two of the three target proteins (R1, BEI, BEII proteins). In this embodiment of the invention, the inhibition of the expression of the third target gene and/or the reduction of the activity of the third target protein takes place by means of a separate foreign nucleic acid molecule which contains the corresponding genetic information for exerting this inhibiting effect.

In another embodiment of the invention, it is not a triple construct that is introduced into the genome of the plant cell but several different foreign nucleic acid molecules, one of these foreign nucleic acid molecules being for example a DNA molecule which, for instance, is a co-suppression construct leading to a reduction of the expression of one or more endogenous R1 genes, and a further foreign nucleic acid molecule being a DNA molecule encoding, for example, an antisense RNA leading to a reduction of the expression of one or more endogenous BEI and/or BEII genes. In principle, as regards the construction of the foreign nucleic acid molecules, it is also suitable to use every combination of antisense, co-suppression and ribozyme constructs or in-vivo mutagenesis, which all lead to a simultaneous reduction of the gene expression of endogenous genes encoding one or more R1, BEI and BEII proteins or which lead to a simultaneous reduction of the activity of one or more R1, BEI and BEII proteins.

In this case, the foreign nucleic acid molecules can be introduced simultaneously ("co-transformation") or consecutively, i.e. one after the other ("super transformation"), into the genome of the plant cell.

In another embodiment of the invention, at least one antisense RNA is expressed for reducing the activity of one or more R1 proteins and/or BEI proteins and/or BEII proteins in plant cells.

For inhibiting the gene expression by means of antisense or co-suppression technology, it is possible to use for instance a DNA molecule which comprises the entire sequence encoding an R1 and/or BEI and/or BEII protein, including flanking sequences that may optionally be present, as well as DNA molecules which only comprise parts of the coding sequence and/or flanking sequences, wherein these parts must be long enough to lead to an antisense effect or a co-suppression effect in the cells. In general, sequences having a minimum length of 15 bp, preferably a length of 100 to 500 bp, in particular sequences having a length of more than 500 bp, are suitable for an efficient antisense or co-suppression inhibition. Usually, DNA molecules which are shorter than 5000 bp, preferably sequences which are shorter than 2500 bp are used.

For antisense or co-suppression approaches, it is also suitable to use DNA sequences which have a high degree of homology to the sequences that occur endogenously in the plant cell and that encode R1, BEI or BEII proteins. The minimum degree of homology should be higher than 65%. It is preferred to use sequences having a homology of at least 90%, in particular between 95 and 100%.

Moreover, also introns, i.e. of non-coding regions of genes encoding R1, BEI and/or BEII proteins, are conceivable for use to achieve an antisense or a co-suppressive effect.

The use of intron sequences for inhibiting the gene expression of genes encoding proteins of the starch biosynthesis has been described in the international patent applications WO 97/04112, WO 97/04113, WO 98/37213, WO 98/37214.

The person skilled in the art knows how to achieve an antisense and a co-suppressive effect. The method of co-suppression inhibition was described, for instance, in Jorgensen (*Trends Biotechnol.* 8 (1990), 340-344), Niebel et al. (*Curr. Top. Microbiol. Immunol.* 197 (1995), 91-103), Flavell et al. (*Curr. Top. Microbiol. Immunol.* 197 (1995), 43-46), Palaqui and Vaucheret (*Plant. Mol. Biol.* 29 (1995), 149-159), Vaucheret et al. (*Mol. Gen. Genet.* 248 (1995), 311-317), de Borne et al. (*Mol. Gen. Genet.* 243 (1994), 613-621).

The skilled persons are also familiar with the expression of ribozymes for reducing the activity of certain enzymes in cells and is described, for example, in EP-B1 0 321 201. The expression of ribozymes in plant cells has been described, for instance, in Feyter et al. (Mol. Gen. Genet. 250, (1996), 329-338).

Furthermore, the R1 and/or BEI and/or BEII activity in plant cells may also be reduced by the so-called "in-vivo mutagenesis" in which a hybrid RNA-DNA oligonucleotide ("chimeroplast") is introduced into cells by means of the transformation of cells (Kipp, P.B. et al., Poster at the 5<sup>th</sup> International Congress of Plant Molecular Biology, 21 - 27 September 1997, Singapore; R.A. Dixon and C.J. Arntzen, Meeting report to "Metabolic Engineering in Transgenic Plants", Keystone Symposia, Copper Mountain, CO, USA, TIBTECH 15, (1997), 441-447; international patent application WO 95/15972; Kren et al., Hepatology 25, (1997), 1462-1468; Cole-Strauss et al., Science 273, (1996), 1386-1389).

A part of the DNA component of the RNA-DNA oligonucleotide is homologous to a nucleic acid sequence of an endogenous R1, BEI and/or BEII gene, it has, however, a mutation in comparison with an endogenous R1, BEI and/or BEII gene or it contains a heterologous region which is embraced by the homologous regions.

It is possible to transfer the mutation or heterologous region contained in the DNA component of the RNA-DNA molecule into the genome of a plant cell by means of base pairing of the homologous regions of the RNA-DNA oligonucleotide and the endogenous nucleic acid molecule, followed by homologous recombination. As a result, the activity of one or more R1, BEI and/or BEII proteins is reduced.

In addition, the reduction of the R1 and/or BEI and/or BEII activity in the plant cells may also be caused by the simultaneous expression of sense and antisense RNA molecules of the target gene that is to be repressed, preferably of the R1 and/or the BEI and/or the BEII gene.

This can be achieved, for instance, by using chimeric constructs which contain inverted repeats of the respective target gene or of parts of the target gene. In this case, the chimeric constructs encode sense and antisense RNA molecules of the respective target gene. Sense and antisense RNA are synthesised simultaneously *in*

*planta* as one RNA molecule, wherein the sense and the antisense RNA may be separated from each other by a spacer and form a double-stranded RNA-molecule. It was possible to show that the introduction of inverted repeat DNA constructs into the genome of plants is a very efficient method for repressing the genes corresponding to the inverted repeat DNA constructs (Waterhouse et al., Proc. Natl. Acad. Sci. USA 95, (1998), 13959-13964; Wang and Waterhouse, Plant Mol. Biol. 43, (2000), 67-82; Singh et al., Biochemical Society Transactions vol. 28, part 6 (2000), 925-927; Liu et al., Biochemical Society Transactions vol. 28, part 6 (2000); 927-929; Smith et al., Nature 407, (2000), 319-320; international patent application WO 99/53050 A1). Sense and antisense sequences of the target gene or of the target genes can also be expressed separately using the same or different promoters (Nap, J.-P. et al., 6<sup>th</sup> International Congress of Plant Molecular Biology, Quebec, 18 – 24 June 2000; Poster S7-27, lecture session S7).

Thus, it is also possible to reduce the R1 and/or BEI and/or BEII activity in the plant cells by the production of double-stranded RNA molecules of R1 and/or BEI and/or BEII genes. Preferably, inverted repeats of DNA molecules of R1 and/or BEI and/or BEII genes or cDNAs are introduced into the genome of plants for this purpose, wherein the DNA molecules to be transcribed (R1, BEI or BEII gene or cDNA or fragments of these genes or cDNAs) are under the control of a promoter which controls the expression of said DNA molecules.

Furthermore, it is known that in *planta* the formation of double-stranded RNA molecules of promoter DNA molecules in plants can lead *in trans* to a methylation and a transcriptional inactivation of homologous copies of these promoters which are called target promoters in the following (Mette et al., EMBO J. 19, (2000), 5194-5201).

Therefore, it is possible to reduce the gene expression of a certain target gene (e.g. R1, BEI or BEII gene) by means of the inactivation of the target promoter, the target gene being naturally controlled by this target promoter.

This means that, in this case, in contrast to the original function of promoters in plants, the DNA molecules which comprise the target promoters of the genes to be

repressed (target genes) are not used as control elements for the expression of genes or cDNAs but as transcribable DNA molecules themselves.

For producing the double-stranded target promoter RNA molecules *in planta* which may be present there as RNA hairpin molecules, constructs are preferred to be used which contain inverted repeats of the target promoter DNA molecule, the target promoter DNA molecules being under the control of a promoter which controls the gene expression of said target promoter DNA molecules.

Then, these constructs are introduced into the genome of plants. The expression of the inverted repeats of said target promoter DNA molecules leads to the formation of double-stranded target promoter RNA molecules *in planta* (Mette et al., EMBO J. 19, (2000), 5194-5201). In this way, it is possible to inactivate the target promoter.

Therefore, the R1 and/or BEI and/or BEII activity in the plant cells can also be reduced by generating double-stranded RNA molecules of promoter sequences of R1 and/or BEI and/or BEII genes. For this purpose, inverted repeats of promoter DNA molecules of R1 and/or BEI and/or BEII promoters are preferred to be introduced into the genome of plants, the target promoter DNA molecules to be transcribed (R1, BEI and/or BEII promoter) being under the control of a promoter which controls the expression of said target promoter DNA molecules. The promoter sequences from R1 and/or BEI and/or BEII genes necessary for carrying out the present embodiment can be provided by methods known to the skilled person and described in the literature such as in Sambrook and Russell (2001), Molecular Cloning, CSH Press, Cold Spring Harbor, NY, USA. The methods may for example include the preparation of a genomic library from the plant in which the activity of R1, BEI and BEII proteins shall be reduced, screening of the library for clones containing the sequence flanking the coding region of the respective gene in 5'-direction by the help of a probe comprising a coding sequence for the R1 or BEI or BEII protein as described above and finally sequencing positive clones by conventional techniques.

Moreover, the skilled person knows that the reduction of activity of one or more R1, BEI and/or BEII proteins can be achieved by means of the expression of non-functional derivatives, in particular trans-dominant mutants of such proteins, and/or by means of the expression of antagonists/inhibitors of such proteins.

Antagonists/inhibitors of such proteins comprise, for instance, antibodies, antibody fragments or molecules having similar binding properties. A cytoplasmatic scFv antibody, for example, was used for modulating the activity of the phytochrome A protein in genetically modified tobacco plants (Owen, Bio/Technology 10 (1992), 790-4; Review: Franken, E., Teuschel, U. and Hain, R., Current Opinion in Biotechnology 8, (1997), 411-416; Whitelam, Trends Plant Sci. 1 (1996), 268-272).

Therefore, a plant cell of the invention is also subject matter of the present invention, wherein said foreign nucleic acid molecule the presence and/or expression of which causes a reduction of R1, BEI and BEII activity in said plant cell is selected from the group consisting of

- a) DNA molecules which encode at least one antisense RNA leading to a reduction of the expression of endogenous genes encoding R1 proteins and/or BEI proteins and/or the BEII proteins, preferably encoding R1, BEI and BEII proteins;
- b) DNA molecules which, through a co-suppression effect, lead to a reduction of the expression of endogenous genes encoding R1 proteins and/or BEI proteins and/or BEII proteins, preferably encoding R1, BEI and BEII proteins;
- c) DNA molecules encoding at least one ribozyme which specifically cleaves transcripts of endogenous genes encoding R1 proteins and/or BEI proteins and/or BEII proteins, preferably encoding R1, BEI and BEII proteins;
- d) nucleic acid molecules which have been introduced by means of *in-vivo* mutagenesis and which lead to a mutation or an insertion of a heterologous sequence in the genes encoding endogenous R1 proteins and/or BEI proteins and/or BEII proteins, preferably encoding R1, BEI and BEII proteins, wherein the mutation or insertion leads to a reduction of the expression of genes encoding R1 proteins and/or BEI proteins and/or BEII proteins, or the synthesis of inactive R1 and/or BEI and/or BEII proteins; and
- e) DNA molecules which simultaneously encode at least one antisense RNA and at least one sense RNA, wherein said antisense RNA and said sense RNA form a double-stranded RNA molecule which leads to a reduction of the expression

of endogenous genes encoding R1 proteins and/or BEI proteins and/or BEII proteins, preferably encoding R1, BEI and BEII proteins.

In another embodiment, the present invention relates to transgenic plant cells which synthesises a modified starch. The transgenic plant cells of the invention synthesise a modified starch which is modified in its physico-chemical properties, in particular the amylose/amylopectin ratio, the phosphate content, the viscosity behaviour, the size of the starch granules and/or the form of the starch granules in comparison with starch synthesised in wild type plants so that it is more suitable for specific purposes of application.

It was surprisingly found that the composition of the starch is modified in the plant cells of the invention in such a way that it has an amylose content of at least 75% and a reduced phosphate content in comparison with starch from plant cells from corresponding wild type plants, so that said starch is more suitable for specific purposes of application.

A plant cell of the invention which contains modified starch having an amylose content of at least 75% and a reduced phosphate content compared to the starch of corresponding plant cells of wild type plants, the cells not being genetically modified, is also subject matter of the present invention.

In the context of the present invention, the amylose content is determined according to the method by Hovenkamp-Hermelink et al. described below in connection with potato starch (Potato Research 31, (1988), 241-246). This method can also be used for isolated starches of other plant species. The person skilled in the art is familiar with methods for isolating starches.

Within the meaning of the present invention, the term "phosphate content" relates to the content of phosphate bound covalently in form of starch phosphate monoesters.

In the context of the present invention, the expression "reduced phosphate content" means that the overall phosphate content of phosphate covalently bound and/or the phosphate content in the C-6 position of the starch synthesised in the plant cells of the invention is reduced by at least 20%, preferably by at least 50%, more preferably by at least 80% in comparison with starch from plant cells of corresponding wild type plants, the cells not being genetically modified.

Within the meaning of the present invention, the term "phosphate content in the C-6 position" is understood to be the content of phosphate groups which are bound to the carbon atom position "6" of the glucose monomers of the starch. In principle, the positions C2, C3 and C6 of the glucose units may be phosphorylated in the starch *in vivo*. In connection with the present invention, the phosphate content in the C-6 position (= C6-P content) can be determined through the determination of the glucose-6-phosphate by means of an optic-enzymatic test (Nielsen et al., Plant Physiol. 105, (1994), 111-117) (see below).

In the context of the present invention, the expression "overall phosphate content" of the starch is understood to be the content of phosphate bound covalently in form of starch phosphate monoesters in the C2, C3 and C6 position of the glucose units. According to the invention, the content of phosphorylated non-glucans such as, e.g. phospholipids, is not included in the term "overall phosphate content". Thus, phosphorylated non-glucans must be separated quantitatively before determining the overall phosphate content. The skilled person knows methods for separating the phosphorylated non-glucans (e.g. phospholipids) from the starch. Methods for determining the overall phosphate content are known to the person skilled in the art and are described below.

In a preferred embodiment of the invention, the plant cells of the invention synthesise a starch which has a phosphate content in the C-6 position of the glucose monomers of up to 15 nmol C6-P mg<sup>-1</sup> starch, in particular of up to 10 nmol C6-P mg<sup>-1</sup> starch, preferably of up to 7 nmol C6-P mg<sup>-1</sup> starch, more preferably of up to 4 nmol C6-P mg<sup>-1</sup> starch.



In another embodiment, the present invention therefore relates to plant cells according to the invention which synthesise a modified starch, wherein the modified starch is characterised in that it has a modified distribution of the side chains. It has been shown that the starch modified in the plant cells of the invention is characterised not only by an increased amylose content and a reduced phosphate content compared to the starch of corresponding wild type plants, but also by a modified distribution of the side chains.

In this embodiment, the term "modified distribution of the side chains" is understood to be an increase in the proportion of short side chains having a DP of 26 to 31 by at least 50%, preferably by at least 100%, more preferably by at least 150% and especially preferred by at least 200% in comparison with the proportion of short side chains having a DP of 26 to 31 of amylopectin from wild type plants. Moreover, the term "modified distribution of the side chains" means an increase of the proportion of short side chains having a DP of 26 to 31, wherein the increase of the proportion of short side chains having a DP of 26 to 31 is not higher than 800%, in particular not higher than 500% compared to the proportion of short side chains having a DP of 26 to 31 of amylopectin from wild type plants. The quantity "DP" means the degree of polymerisation.

The proportion of short side chains is determined by the determination of the proportion in percent that a certain side chain has in the overall proportion of all side chains. The overall proportion of all side chains is determined through the determination of the overall height of the peaks which represent the polymerisation degrees of DP 6 to 40 in the HPLC chromatogram. The proportion in percent that a certain side chain has in the overall proportion of all side chains is determined by the determination of the ratio of the height of the peak representing this side chain in the HPLC chromatogram to the overall height. The program Chromeleon 6.20 by Dionex, USA can, for instance, be used for determining the peak areas.

In another preferred embodiment, the present invention relates to plants cells of the invention which synthesise a modified starch which form a gel after pasting in a 60%

(w/v)  $\text{CaCl}_2$  solution, the gel having an increased gel strength compared to the gel from starch of corresponding wild type plant cells that have not been genetically modified.

Within the meaning of the present invention, the term "increased gel strength" means an increase in the gel strength by at least 1000%, in particular by at least 2500%, preferably by at least 5000% and more preferably by at least 10,000%, by 40,000% at the most or by 30,000% at the most in comparison with the gel strength of starch of corresponding wild type plant cells that have not been genetically modified.

In the context of the present invention, the gel strength is to be determined by means of a texture analyzer under the conditions described below. In this case, the pasting of the starch is achieved in an aqueous 60% (w/v)  $\text{CaCl}_2$  solution since in a purely aqueous system, it is not possible to achieve pasting of the starch at normal pressure.

In a further preferred embodiment, the present invention relates to plant cells of the invention which, in addition to the aforementioned properties, the starch of which has a modified morphology of the starch granules.

In comparison with high-amylose starches which are known so far, in particular with high-amylose potato starches, the starches of the plant cells of the invention are not only modified in the amylose content, the phosphate content, the distribution of the side chains, the viscosity behaviour and the gel formation behaviour, but also in a modified morphology of the starch granules, which renders these starches more suitable for certain purposes of application.

These starches, in particular the potato starches, could, for instance, be used instead of rice starches since, after mechanical fragmentation, the starches of the invention have an average size of the starch granules which is similar to that of rice starches. Compared to rice starches, the starches of the invention, in particular the potato starches, however have the advantage that they can be sedimented to larger units having the form of a bunch of grapes (cf. Example 2) as small starch granules form bunch-of-grapes-like agglomerations, which may be of advantage in the extraction and processing of the starch and by which the costs may be reduced.

Preferably, the morphology of the starch granules contained in the plant cells of the invention is characterised by an agglomeration of small starch granules having the form of a bunch of grapes.

In a preferred embodiment, the starches contained in the plant cells of the invention are characterised in that the average granule size is reduced compared to the average granule size of corresponding cells of wild type plants which are not genetically modified.

In the context of the present invention, the term "average granule size" means the granule size which can be determined using, for instance, a photo sedimentometer of the type "Lumosed FS1" by Retsch GmbH (see below).

In a further preferred embodiment of the invention, a reduced average granule size is a reduction of the average granule size by at least 20%, preferably by at least 40% and more preferably by at least 60%.

In another preferred embodiment, the starches of the plant cells of the invention are characterised by an average granule size of less than 20  $\mu\text{m}$ , in particular of less than 18  $\mu\text{m}$ , preferably of less than 16  $\mu\text{m}$  and more preferably of 10 – 15  $\mu\text{m}$ .

In another preferred embodiment of the invention, the starches of the plant cells of the invention are characterised in that the proportion of granules having an average granule size of less than 20  $\mu\text{m}$  is at least 70%, preferably at least 75% and more preferably at least 80%.

After mechanical fragmentation of the starch, which may be carried out as described below, the starches of the plant cells of the invention have a proportion of granules having a granule size of less than 20  $\mu\text{m}$  of at least 80%, preferably of at least 90% and more preferably of at least 95%.

A plurality of techniques is available for introducing DNA into a plant host cell. These techniques comprise the transformation of plant cells with T-DNA using *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes* as a transformation means, the fusion of protoplasts, injection, the electroporation of DNA, the introduction of DNA by means of the biolistic approach, and other possibilities.

The use of the *Agrobacteria*-mediated transformation of plant cells has been researched into intensively and described sufficiently in EP120516; Hoekema, in: The Binary Plant Vector System, Offsetdrukkerij Kanters B.V., Alblisserdam (1985), Chapter V; Fraley et al., Crit. Rev. Plant Sci. 4, 1-46 and An et al. EMBO J. 4, (1985), 277-287. For the transformation of potato, see, for example, Rocha-Sosa et al. (EMBO J. 8, (1989), 29-33.).

The transformation of monocotyledonous plants by means of *Agrobacterium*-based vectors has also been described (Chan et al., Plant Mol. Biol. 22, (1993), 491-506; Hiei et al., Plant J. 6, (1994) 271-282; Deng et al., Science in China 33, (1990), 28-34; Wilmink et al., Plant Cell Reports 11, (1992), 76-80; May et al., Bio/Technology 13, (1995), 486-492; Conner and Domisse, Int. J. Plant Sci. 153 (1992), 550-555; Ritchie et al., Transgenic Res. 2, (1993), 252-265). An alternative system for the transformation of monocotyledonous plants is the transformation by the biolistic approach (Wan and Lemaux, Plant Physiol. 104, (1994), 37-48; Vasil et al., Bio/Technology 11 (1993), 1553-1558; Ritala et al., Plant Mol. Biol. 24, (1994), 317-325; Spencer et al., Theor. Appl. Genet. 79, (1990), 625-631), protoplast transformation, the electroporation of partially permeabilised cells, and the introduction of DNA by means of glass fibers. The transformation of maize, in particular, has been described repeatedly in the literature (cf., for example, WO 95/06128, EP0513849, EP0465875, EP0292435; Fromm et al., Biotechnology 8, (1990), 833-844; Gordon-Kamm et al., Plant Cell 2, (1990), 603-618; Koziel et al., Biotechnology 11 (1993), 194-200; Moroc et al., Theor. Appl. Genet. 80, (1990), 721-726).

The successful transformation of other cereal species has also been described, for example in the case of barley (Wan and Lemaux, see above; Ritala et al., see above; Krens et al., Nature 296, (1982), 72-74) and wheat (Nehra et al., Plant J. 5, (1994), 285-297). For the expression of the foreign nucleic acid molecule (foreign

nucleic acid molecules), in principle, any promoter which is active in plant cells can be used. The promoter can be chosen in such a way that expression in the plants according to the invention is constitutive, or only in a particular tissue, at a particular point in time of plant development, or at a point in time determined by external factors. With respect to the plant, the promoter may be homologous or heterologous. Examples of suitable promoters are the promoter of the cauliflower mosaic virus 35S RNA and the ubiquitin promoter from maize for constitutive expression, the patatin gene promoter B33 (Rocha-Sosa et al., EMBO J. 8 (1989), 23-29), the MCPI promoter of the metallocarboxypeptidase inhibitor gene from potato (Hungarian patent application HU9801674) or the GBSSI promoter from potato (international patent application WO 92/11376) for tuber-specific expression in potatoes, or a promoter which ensures expression only in photosynthetically active tissues, for example the ST-LS1 promoter (Stockhaus et al., Proc. Natl. Acad. Sci. USA 84 (1987), 7943-7947; Stockhaus et al., EMBO J. 8 (1989), 2445-2451), the Ca/b promoter (see, for example, US 5656496, US 5639952, Bansal et al., Proc. Natl. Acad. Sci. USA 89, (1992), 3654-3658) and the rubisco SSU promoter (see, for example, US 5034322, US 4962028), or the glutelin promoter for an endosperm-specific expression (Leisy et al., Plant Mol. Biol. 14 (1990), 41-50; Zheng et al., Plant J. 4, (1993), 357-366; Yoshihara et al., FEBS Lett. 383, (1996), 213-218), the shrunken-1 promoter (Werr et al., EMBO J. 4, (1985), 1373-1380), the HMG promoter from wheat, the USP promoter, the phaseolin promoter, or promoters of maize zein genes (Pedersen et al., Cell 29, (1982), 1015-1026; Quatroccio et al., Plant Mol. Biol. 15 (1990), 81-93). The expression of the foreign nucleic acid molecule (the foreign nucleic acid molecules) is of particular advantage in organs of the plant that store starch. Such organs are, e.g., the tuber of the potato plant or the kernels or the endosperm of maize, wheat or rice plants. Thus, promoters mediating the expression in these organs are preferred to be used.

However, it is also possible to use promoters which are only activated at a point in time which is determined by external factors (see, for example, WO 93/07279). Promoters of heat shock proteins, which permit simple induction, may be of particular interest in this context. Furthermore, seed-specific promoters such as, for example, the *Vicia faba* USP promoter which ensures seed-specific expression in *Vicia faba*

and other plants (Fiedler et al., Plant Mol. Biol. 22, (1993), 669-679; Bäumlein et al., Mol. Gen. Genet. 225, (1991), 459-467) can be used. Other promoters which can be employed are fruit-specific promoters as described, for example, in WO 91/01373.

A termination sequence which serves for the correct termination of the transcription and for adding a poly-A tail to the transcript, which is understood to have a function in stabilising the transcripts, may furthermore be present. Such elements have been described in the literature (cf., for example, Gielen et al., EMBO J. 8 (1989), 23-29) and are freely exchangeable.

The plant cells according to the invention may belong to any plant species, i.e. to monocotyledonous or dicotyledonous plants. They are preferably plant cells from agriculturally useful plants, i.e. plants which are grown by man for the purposes of nutrition or for technical, in particular industrial, purposes. The invention preferably relates to fibre-forming (for example flax, hemp, cotton), oil-storing (for example rape, sunflower, soy bean), sugar-storing (for example sugar beet, sugar cane, sugar millet) and protein-storing plants (for example leguminous plants).

In a further preferred embodiment, the invention relates to fodder plants, in particular forage grass and pasture grass (alfalfa, clover, etc.) and vegetable plants (for example tomato, lettuce, chicory).

In another preferred embodiment, the invention relates to plant cells from starch-storing plants (for example wheat, barley, oat, rye, potato, maize, rice, pea, cassava), particularly preferred are plant cells from potato.

The plant cells of the invention can be used for regenerating whole plants.

The plants obtainable by regenerating the transgenic plant cells of the invention are also subject matter of the present invention.

Furthermore, plants which contain the transgenic plant cells of the invention are also subject matter of the invention.

The transgenic plants may, in principle be plants belonging to any plant species, i.e.

both monocotyledonous and dicotyledonous plants. They are preferably useful plants, i.e. plants which are grown by man for the purposes of nutrition or for technical, in particular industrial, purposes. The invention preferably relates to plant cells of fibre-forming (for example flax, hemp, cotton), oil-storing (for example rape, sunflower, soy bean), sugar-storing (for example sugar beet, sugar cane, sugar millet) and protein-storing plants (for example leguminous plants). In a further preferred embodiment, the invention relates to fodder plants, in particular forage grass and pasture grass (alfalfa, clover, etc.) and vegetable plants (for example tomato, lettuce, chicory).

In another preferred embodiment, the invention relates to starch-storing plants (for example wheat, barley, oat, rye, potato, maize, rice, pea, cassava), particularly preferred are potato plants.

The present invention also relates to a method for the production of a transgenic plant cell which synthesises a modified starch, wherein a plant cell is genetically modified by introducing one or more foreign nucleic acid molecules, the presence and/or expression of which leads to a reduction of the activity of R1, BEI and BEII proteins compared to corresponding plants cells of wild type plants, the cells not being genetically modified.

In a preferred embodiment of the method of the invention, the modified starch is characterised in that it has an amylose content of at least 75% and a reduced phosphate content in comparison with starch from corresponding wild type plants which are not genetically modified.

The present invention also relates to a method for producing a transgenic plant which synthesises modified starch, wherein

- a) a plant cell is genetically modified by introducing one or more foreign nucleic acid molecules the presence and/or expression of which leads to a reduction of the activity of R1, BEI and BEII proteins compared to corresponding plant cells of wild type plants, the cells not being genetically modified;
- b) a plant is regenerated from the cell produced according to a); and
- c) optionally further plants are produced from the plant produced according to step

b).

In a preferred embodiment of the method of the invention, the modified starch is characterised in that it has an amylose content of at least 75% and a reduced phosphate content compared to the starch from corresponding wild type plants which are not genetically modified.

In another embodiment of the method of the invention, the modified starches moreover have a modified distribution of the side chains and/or a modified morphology of the starch granules and/or a reduced average size of the starch granules and/or form a gel after pasting in an aqueous 60% (w/v)  $\text{CaCl}_2$  solution, the gel having an increased gel strength in comparison with a gel of starch from corresponding wild type plants which are not genetically modified.

The same as has already been said above in connection with the plant cells of the invention also applies to the genetic modification introduced according to step a). Regeneration of plants according to step b) can be made using methods known to the person skilled in the art.

Further plants of the methods of the invention can be produced according to step c) by means of vegetative propagation (for example using cuttings, tubers or by means of callus culture and regeneration of whole plants) or by generative propagation. Generative propagation is preferably done under controlled conditions, i.e. selected plants having specific properties are crossed with each other and propagated. The person skilled in the art obviously knows that, for producing the plant cells and plants of the invention, also transgenic plants can be used in which the activity of one or two of the aforementioned proteins has already been reduced and which, according to the method of the invention, only have to be genetically modified in such a way that the activity of the second or third protein is also reduced.

In addition, the skilled person knows that the aforementioned super-transformation is not necessarily carried out with primary transformants but preferably with pre-selected stable transgenic plants which advantageously have already been tested for, e.g. fertility, stable expression of the foreign gene, hemizygoty and heterozygoty, etc. in corresponding experiments.



The present invention also relates to the plants obtainable by the methods of the invention.

The present invention also relates to propagation material of plants of the invention containing plant cells of the invention as well as of the plants produced according to the methods of the invention. Within the meaning of the present invention, the term "propagation material" comprises parts of the plant which are suitable for producing progeny by the vegetative or generative route. Examples which are suitable for vegetative propagation are cuttings, callus cultures, rhizomes or tubers. Other propagation material encompasses, for example, fruits, seeds, seedlings, protoplasts, cell cultures and the like. The propagation material is preferably seeds.

Furthermore, the present invention relates to the use of one or more foreign nucleic acid molecules encoding proteins having the enzymatic activity of R1, BEI and BEII proteins and to the use of fragments of said foreign nucleic acid molecules for producing plant cells or plants of the invention synthesising a modified starch.

In another embodiment of the invention, the plant cells of the invention synthesise a modified starch due to the use according to the invention of one or more foreign nucleic acid molecules, the modified starch being characterised in that it has an amylose content of at least 75% and/or a reduced phosphate content compared to the starch of corresponding wild type plants which have not been genetically modified and/or a modified distribution of the side chains and/or a modified morphology of the starch granules and/or a reduced average size of the starch granules and/or a modified starch which forms a gel after pasting in an aqueous 60% (w/v)  $\text{CaCl}_2$  solution, the gel having an increased gel strength in comparison with a gel of starch from corresponding wild type plant cells which are not genetically modified.

In another embodiment, the present invention relates to the use of one or more foreign nucleic acid molecules for producing plants of the invention, wherein the

foreign nucleic acid molecule is a molecule, or the foreign nucleic acid molecules are several molecules selected from the group consisting of

- a) DNA molecules which encode at least one antisense RNA leading to a reduction of the expression of endogenous genes encoding the R1 proteins and/or BEI proteins and/or the BEII proteins, preferably encoding R1, BEI and BEII proteins;
- b) DNA molecules which, through a co-suppression effect, lead to the reduction of the expression of endogenous genes encoding R1 proteins and/or BEI proteins and/or BEII proteins, preferably encoding R1, BEI and BEII proteins;
- c) DNA molecules encoding at least one ribozyme which specifically cleaves transcripts of endogenous genes encoding R1 proteins and/or BEI proteins and/or BEII proteins, preferably encoding R1, BEI and BEII proteins;
- d) nucleic acid molecules which have been introduced by means of *in-vivo* mutagenesis and which lead to a mutation or an insertion of a heterologous sequence in the genes encoding endogenous R1 proteins and/or BEI proteins and/or BEII proteins, preferably encoding R1, BEI and BEII proteins, wherein the mutation or insertion leads to a reduction of the expression of the genes encoding R1 proteins and/or BEI proteins and/or BEII proteins, or the synthesis of inactive R1 and/or BEI and/or BEII proteins; and
- e) DNA molecules which simultaneously encode at least one antisense RNA and at least one sense RNA, wherein said antisense RNA and said sense RNA form a double-stranded RNA molecule which leads to a reduction of the expression of endogenous genes encoding R1 proteins and/or BEI proteins and/or BEII proteins, preferably encoding R1, BEI and BEII proteins.

As has already been explained before, the foreign nucleic acid molecules can be introduced simultaneously or consecutively, i.e. one after the other, into the genome of the plant cell. The simultaneous introduction of the foreign nucleic acid molecules saves time and costs, i.e. the co-transformation in which, preferably in one transformation experiment according to the aforementioned methods of the invention, foreign nucleic acid molecules are introduced into the plant cell, the presence and optionally the expression of which lead to the reduction of the activity

of one or more R1 proteins occurring endogenously in the plant cell and to the reduction of the activity of one or more BEI proteins occurring endogenously in the plant cell and to the reduction of the activity of one or more BEII proteins occurring endogenously in the plant cell in comparison with corresponding plant cells of wild type plants, the cells not being genetically modified.

Thus, the present invention also relates to compositions containing at least one of the foreign nucleic acid molecules defined according to the invention, these foreign nucleic acid molecules being suitable for producing the transgenic plant cells and/or the transgenic plants of the invention. Preferably, the presence and/or expression of these foreign nucleic acid molecules in plant cells leads to the reduction of the activity of R1 and BEI and BEII proteins compared to corresponding plant cells of wild type plants, the cells not being genetically modified.

In this case, in the composition of the invention, the nucleic acid molecules the presence and/or expression of which in the plant cell and/or the plant leads to the reduction of the activity of R1 and BEI and BEII proteins compared to corresponding plant cells of wild type plants, the cells not being genetically modified, can be contained either separately or together in one recombinant nucleic acid molecule. In the former case, the composition of the invention can, for instance, contain two or more recombinant nucleic acid molecules and/or vectors the joint presence of which in the plant cell leads to said phenotype. In the latter case, a recombinant nucleic acid molecule contains the genetic information leading to the reduction of the activity of R1 and BEI and BEII proteins compared to corresponding plant cells of wild type plants, the cells not being genetically modified.

In such a recombinant molecule, for instance, the above-described foreign nucleic acid molecules the presence and/or expression of which in a plant cell leads to the reduction of the activity of R1 and BEI and BEII proteins compared to corresponding plant cells of wild type plants, the cells not being genetically modified, can be present as one chimeric gene or as separate genes. Examples of such double or multiple constructs have been described numerously in the literature.

The aforementioned recombinant nucleic acid molecules can be present in any host

cell.

In another embodiment, the present invention therefore also relates to a host cell, in particular a plant cell, containing a composition of the invention.

The plant cells and plants of the invention synthesise a starch, in particular in their starch-storing organs, which is modified in its physico-chemical properties, in particular the phosphate content and/or the amylose content, preferably the phosphate content and the amylose content, and/or the distribution of the side chains and/or the viscosity behaviour and/or the morphology of the starch granules and/or the average size of the starch granules in comparison with starch synthesised in wild type plants.

Thus, starch which is obtainable from the plant cells, plants and/or propagation material of the invention is also subject matter of the invention.

In a preferred embodiment, the starch of the invention is characterised in that it has an amylose content of at least 75% and a reduced phosphate content in comparison with starch from corresponding wild type plants which are not genetically modified.

The meaning of the term "increased gel strength" has already been defined in connection with the description of the plant cells of the invention.

In comparison with high-amylose starches which are known so far, in particular with high-amylose potato starches, the starches of the invention are not only modified in the amylose content, the phosphate content, the distribution of the side chains, the viscosity behaviour and the gel formation behaviour, but also in a modified morphology of the starch granules, which renders these starches more suitable for certain purposes of application.

The starches of the invention, in particular the potato starches, could, for instance, be used instead of rice starches since, after mechanical fragmentation, the starches of the invention have an average size of the starch granules which is similar to that of rice starches. Compared to rice starches, the starches of the invention, in particular

the potato starches, however have the advantage that they can be sedimented to larger units having the form of a bunch of grapes (cf. Example 2) as small starch granules form bunch-of-grapes-like agglomerations, which may be of advantage in the extraction and processing of the starch and by which the costs may be reduced.

Preferably, morphology of the starch granules of the starch of the invention is characterised by an agglomeration of small starch granules having the form of a bunch of grapes.

In another embodiment, the starches of the invention are characterised in that the average granule size is reduced compared to the average granule size of corresponding wild type plants which are not genetically modified.

In the context of the present invention, the term "average granule size" means the granule size which can be determined using, for instance, a photo sedimentometer of the type "Lumosed FS1" by Retsch GmbH (see below).

In another embodiment of the invention, a reduced average granule size is a reduction of the average granule size by at least 20%, preferably by at least 40% and more preferably by at least 60%.

In another embodiment, the starches of the invention are characterised by an average granule size of less than 20  $\mu\text{m}$ , in particular of less than 18  $\mu\text{m}$ , preferably of less than 16  $\mu\text{m}$  and more preferably of 10 – 15  $\mu\text{m}$ .

In another embodiment of the invention, the starches of the invention are characterised in that the proportion of granules having an average granule size of less than 20  $\mu\text{m}$  is at least 70%, preferably at least 75% and more preferably at least 80%.

After mechanical fragmentation of the starch, which may be carried out as described below, the starches of the invention have a proportion of granules having a granule

size of less than 20  $\mu\text{m}$  of at least 80%, preferably of at least 90% and more preferably of at least 95%.

In a particularly preferred embodiment, the starch of the invention is a potato starch.

Moreover, the present invention relates to a method for producing the starches of the invention comprising the step of extracting the starch from a plant (cell) of the invention and/or from starch-storing parts of such a plant.

Preferably, such a method also comprises the step of harvesting the cultivated plants and/or starch-storing parts of said plants prior to extracting the starch and, particularly preferably, the step of cultivating the plants of the invention prior to the harvesting.

The person skilled in the art knows methods for extracting the starch from plants or from starch-storing parts of plants. Furthermore, methods for extracting the starch from various starch-storing plants have been described, e.g. in "Starch: Chemistry and Technology (editors.: Whistler, BeMiller and Paschall (1994), 2<sup>nd</sup> edition, Academic Press Inc. London Ltd.; ISBN 0-12-746270-8; cf., e.g. chapter XII, page 412-468: maize and sorghum starches: production; by Watson; Chapter XIII, page 469-479: tapioca, arrowroot and sago starches: production; by Corbishley and Miller; Chapter XIV, page 479-490: potato starch: production and uses; by Mitch; Chapter XV, page 491 to 506: wheat starch: production, modification and uses; by Knight and Oson; and Chapter XVI, page 507 to 528: rice starch: production and uses; by Rohmer and Klem; maize starch: Eckhoff et al., Cereal Chem. 73 (1996) 54-57), the extraction of maize starch on an industrial scale is generally achieved by wet milling. Apparatuses usually used in processes for extracting starch from plant materials are separators, decanters, hydrocyclones, spray dryers and fluidized-bed dryers.

Moreover, starch which is obtainable using the aforementioned method of the invention is also subject matter of the invention.

The starches according to the invention can be modified afterwards by processes known to the skilled person and are suitable, in their unmodified or modified forms,

known to the skilled person and are suitable, in their unmodified or modified forms, for a variety of applications in the food or non-food sector.

The figures show:

**Figure 1:** Schematic representation of the expression vector ME 5/6 as described further below.

**Figure 2:** Light-microscopic view of starch granules of wild type potato plants.

**Figure 3:** Light-microscopic view of starch granules of 072VL036 potato plants having a reduced gene expression of the R1 and BEI gene.

**Figure 4:** Light-microscopic view of starch granules of 203MH010 potato plants according to the invention.

The following methods were used in the Examples:

Analysis of the starch

a) Determination of the amylose/amylopectin ratio

Starch was isolated from potato plants according to standard techniques and the ratio of amylose to amylopectin was determined using the method described by Hovenkamp-Hermelink et al. (Potato Research 31, (1988), 241-246).

b) Determination of the phosphate content

The positions C2, C3 and C6 of the glucose units may be phosphorylated in the starch. For determining the C6-P content of the starch, 50 mg starch are hydrolysed in 500  $\mu$ l 0.7 M HCl for 4 hours at 95°C. Then, the mixtures are centrifuged for 10 min at 15,500 g and the supernatants are taken. 7  $\mu$ l of the supernatants are mixed with 193  $\mu$ l imidazole buffer (100 mM imidazole, pH 7.4; 5 mM MgCl<sub>2</sub>, 1 mM EDTA and 0.4 mM NAD). The measuring was carried out in the photometer at 340 nm. After establishing a basic absorption, the enzyme reaction was started by adding 2 u glucose-6 phosphate dehydrogenase (from *Leuconostoc mesenteroides*, Boehringer Mannheim). The change in the absorption is directly proportional to the concentration of the G-6-P content of the starch.

The overall phosphate content was determined according to the method by Ames (Methods in Enzymology VIII, (1966), 115-118).

30  $\mu$ l ethanolic magnesium nitrate solution are added to about 50 mg starch and ashed for three hours at 500°C in a muffle furnace. 300  $\mu$ l 0.5 M hydrochloric acid are added to the residue and incubated for 30 min at 60°C. Then, an aliquot is filled to 300  $\mu$ l 0.5 M hydrochloric acid, added to a mixture of 100  $\mu$ l 10% ascorbic acid and 600  $\mu$ l 0.42% ammonium molybdate in 2 M sulphuric acid and incubated for 20 min at 45°C.



Then, a photometric measurement is conducted at 820 nm, using a phosphate calibration series as a standard.

c) Determination of the gel strength (texture analyser)

2 g starch (TS) are dissolved in 25 ml of an aqueous 60% (w/v)  $\text{CaCl}_2$  solution and pasting is achieved in an RVA apparatus (temperature program: cf. item d) "Determination of the viscosity properties by means of a Rapid Visco Analyser (RVA)") and then it is stored in a closed container for 24 hours at room temperature. The samples are fixed under a probe (cylindrical stamp with a planar surface) of a texture analyser TA-XT2 by Stable Micro Systems (Surrey, UK) and the gel strength is determined using the following parameters:

- test speed                      0.5 mm/s
- depth of penetration        7 mm
- contact area                  113 mm<sup>2</sup>
- pressure                        2 g

d) Determination of the viscosity properties by means of a Rapid Visco Analyser (RVA)

2 g starch (TS) are added to 25 ml  $\text{H}_2\text{O}$  and used for the analysis in a Rapid Visco Analyser (Newport Scientific Pty Ltd., Investment Support Group, Warriewood NSW 2102, Australia). The apparatus is used according to the manufacturer's instructions. For determining the viscosity of the aqueous solution of the starch, the starch suspension is first heated to 50°C for 1 min, then it is heated from 50°C to 95°C at a speed of 12°C per minute. Subsequently, the temperature is maintained at 95°C for 2.5 min. Then, the solution is cooled down from 95°C to 50°C at a speed of 12°C per minute. The viscosity is determined over the whole time.

e) Determination of glucose, fructose and sucrose

The content of glucose, fructose and sucrose is determined according to the method described by Stitt et al. (Methods in Enzymology 174, (1989), 518-552).

f) Analysis of the distribution of the side chains of the amylopectin by means of ion exchange chromatography

For separating amylose from amylopectin, 200 mg starch are dissolved in 50 ml-reaction vessels with 12 ml 90% (v/v) DMSO in H<sub>2</sub>O. After adding 3 volumes ethanol, the precipitate is separated by a 10 min-centrifugation at about 1800 g at room temperature (RT). The pellet is then washed with 30 ml ethanol, dried and dissolved in 40 ml 1% (w/v) NaCl solution at 75°C. After cooling down the solution to 30°C, about 90 mg thymol are added slowly and this solution is incubated for at least 60 h at 30°C. Then, the solution is centrifuged for 30 min at 2000 g (RT). 3 volumes ethanol are then added to the supernatant and the precipitating amylopectin is separated by means of 5 min-centrifugation at 2000 g (RT). The pellet (amylopectin) is then washed with ethanol and dried using acetone. By adding DMSO to the pellet, a 1%-solution is prepared 200 µl of which are added to 345 µl water, 10 µl 0.5 M sodium acetate (pH 3.5) and 5 µl isoamylase (dilution of 1:10; Megazyme) and incubated for about 16 h at 37°C. An aqueous 1:5 dilution of this digestion is then filtered with an 0.2 µm-filter and 100 µl of the filtrate are analysed by means of ion exchange chromatography (HPAEC-PAD, Dionex). The separation is carried out with a PA-100 column (with a corresponding pre-column), the detection is carried out amperometrically. The elution conditions are as follows:

solution A - 0.15 M NaOH

solution B – 1 M sodium acetate in 0.15 M NaOH

t (min)	solution A (%)	solution B (%)
5	0	100
35	30	70
45	32	68
60	100	0
70	100	0
72	0	100
80	0	100
stop		

The relative proportion of short side chains in the overall proportion of all side chains is determined by determining the proportion in percent that a certain side chain has in the overall proportion of all side chains. The overall proportion of all side chains is determined through the determination of the overall height of the peaks which represent the polymerisation degrees of DP 6 to 40 in the HPLC chromatogram. The proportion in percent that a certain side chain has in the overall proportion of all side chains is determined by the determination of the ratio of the height of the peak representing this side chain in the HPLC chromatogram to the overall height of all peaks having a DP of 6 to 40. The program Chromeleon 6.20 by Dionex, USA was used for determining the peak heights. The parameters of the evaluation software that were to be adjusted were as follows:

retention time (min)	parameter name	parameter value	channels
0.000	Inhibit Integration	on	All channels
20.000	Lock Baseline	on	All channels
20.600	Inhibit Integration	off	All channels
20.600	Minimum Height	0.001 (Signal)	All channels
45.000	Inhibit Integration	on	All channels

g) Determination of the granule size

Starch was extracted from the potato tubers according to standard methods and washed several times with water in a 10 l-bucket (ratio height of the bucket/diameter of the bucket = approx. 1.1). For obtaining the starch samples which were finally subjected to the determination of the granule size, the starches were left to stand for about 4 h after washing to achieve as complete a sedimentation of the starches as possible.

The granule size was then determined by means of a photo sedimentometer of the type "Lumosed FS1" by Retsch GmbH, Germany using the software V.2.3. The software adjustments were as follows:

data of the substance:	calibration no.	0
	density [kg/m <sup>3</sup> ]	1500
sedimentation fluid:	type	water
	viscosity [Pa s]	0.001
	density [kg/m <sup>3</sup> ]	1000
	addition	-
	measurement data	5 min
	sieve diameter [µm]	250
	passage [%]	100
	measurement range	4.34–117.39 µm
	calibration	N
	temperature	20°C

The distribution of the granule size was determined in an aqueous solution and according to the manufacturer's instructions and based on the literature of e.g. H. Pitsch, Korngrößenbestimmung; LABO-1988/3 Fachzeitschrift für Labortechnik, Darmstadt.

h) Mechanical fragmentation of the starch

About 0.5 g of each starch were placed in a coffee mill (manufacturer: Mellert, type: M85, Germany) and ground six times for 30 s each. Between two intervals, the grinding was interrupted for 20 s each. The distribution of the

44.

granule size was determined as described in item g).

i) Water binding capacity

For determining the water binding capacity (WBC), the supernatant was weighed after separating the soluble portion by centrifugation of the starch swollen at 70°C. The water binding capacity (WBC) of the starch was set in relation to the weighed portion of the starch corrected by the soluble mass.

$$\text{WBC (g/g)} = (\text{residue} - (\text{weighed portion} - \text{soluble portion})) / (\text{weighed portion} - \text{soluble portion}).$$

The expression vector ME5/6 (cf. Fig. 1) was used in the Examples:

Preparation of the expression vector ME5/6

pGSV71 is a derivative of the plasmid pGSV7 which is derived from the intermediary vector pGSV1. pGSV1 is a derivative of pGSC1700 the construction of which has been described by Cornelissen and Vanderwiele (Nucleic Acids Research 17, (1989), 19-25). pGSV1 was obtained from pGSC1700 by deletion of the carbenicillin resistance gene as well as deletion of the T-DNA sequences of the TL-DNA region of the plasmid pTiB6S3.

pGSV7 contains the replication origin of the plasmid pBR322 (Bolivar et al., Gene 2, (1977), 95-113) as well as the replication origin of the *Pseudomonas* plasmid pSV1 (Itoh et al., Plasmid 11, (1984), 206). pGSV7 additionally contains the selectable marker gene *aadA* from the transposon Tn1331 from *Klebsiella pneumoniae* which confers resistance to the antibiotics spectinomycin and streptomycin (Tolmasky, Plasmid 24 (3), (1990), 218-226; Tolmasky and Crosa, Plasmid 29 (1), (1993), 31-40).

The plasmid pGSV71 was obtained by cloning a chimeric *bar* gene between the border regions of pGSV7. The chimeric *bar* gene contains the promoter sequence of the cauliflower mosaic virus for initiating the transcription (Odell et al., Nature 313, (1985), 180), the *bar* gene from *Streptomyces hygrosopicus* (Thompson et al., EMBO J. 6, (1987), 2519-2523) and the 3'-non-translated region of the nopaline

synthase gene of the T-DNA of pTiT37, for terminating the transcription and polyadenylation. The *bar* gene confers resistance to the herbicide glufosinate ammonium.

In position 198-222, the T-DNA contains the right border sequence of the TL-DNA of the plasmid pTiB6S3 (Gielen et al., EMBO J. 3, (1984), 835-846). There is a polylinker sequence between the nucleotide 223-249. The nucleotides 250-1634 contain the P35S3 promoter region of the cauliflower mosaic virus (Odell et al., cf. above). The coding sequence of the phosphinothricine resistance gene (*bar*) from *Streptomyces hygroscopicus* (Thompson et al., 1987, cf. above) is contained between the nucleotides 1635-2186. The two terminal codons at the 5' end of the *bar* wild type gene were replaced by the codons ATG and GAC. There is a polylinker sequence between the nucleotides 2187-2205. The 260 bp-*TaqI* fragment of the non-translated 3' end of the nopaline synthase gene (3'nos) from the T-DNA of the plasmid pTiT37 (Depicker et al., J. Mol. Appl. Genet. 1, (1982), 561-573) is located between the nucleotides 2206 and 2465. The nucleotides 2466-2519 contain a polylinker sequence. The left border sequence of the TL-DNA from pTiB6S3 (Gielen et al., EMBO J. 3, (1984), 835-846) is located between the nucleotides 2520-2544.

The vector pGSV71 was then cleaved with the enzyme *PstI* and blunted. The promoter B33 and the *ocs* cassette were cleaved from the vector pB33-Kan as an *EcoRI-HindIII* fragment, blunted and inserted into the vector pGSV71 which had been cleaved with *PstI* and blunted. The vector obtained served as a starting vector for the construction of ME5/6. An oligonucleotide containing the cleavage sites *EcoRI*, *PacI*, *SpeI*, *SrfI*, *SpeI*, *NofI*, *PacI* and *EcoRI* was inserted into the *PstI* cleavage site between the B33 promoter and the *ocs* element of the vector ME4/6 by duplicating the *PstI* cleavage site. The expression vector obtained was called ME5/6.

#### Description of the vector pSK-Pac:

pSK-Pac is a derivative of pSK Bluescript (Stratagene, USA) into which *PacI* cleavage sites flanking the multiple cloning site (MCS) were inserted.

The following Examples illustrate the invention:

### Example 1

#### **Production of transgenic potato plants having a reduced gene expression of an R1, BEI and BEII gene**

For producing transgenic plants having a reduced activity of a BEI, R1 and BEII protein, first transgenic plants were generated in which the BEI activity and the amount of protein R1 were reduced. For this purpose, both the T-DNA of the plasmid pB33-aR1-Hyg and the T-DNA of the plasmid pB33-a-BE1-Kan were transferred simultaneously into potato plants using *Agrobacteria* as described by Rocha-Sosa et al. (EMBO J. 8, (1989), 23-29).

For constructing the plasmid pB33-aR1-Hyg and the plasmid pB33-aBE1-Kan, first the expression vectors pB33-Kan and pb33-Hyg, respectively, were constructed. For this purpose, the promoter of the patatin gene B33 from *Solanum tuberosum* (Rocha-Sosa et al., 1989, cf. above) was ligated as *DraI* fragment (nucleotides -1512 to +14) into the vector pUC19 (GenBank Acc. No. M77789) which had been cleaved with *SstI*, the ends of said vector having been blunted by means of the T4-DNA polymerase. In this way, the plasmid pUC19-B33 was obtained. The B33 promoter was cleaved from this plasmid with *EcoRI* and *SmaI* and ligated into the vector pBinAR which had been cleaved correspondingly. In this way, the plant expression vector pB33-Kan was obtained. The plasmid pBinAR is a derivative of the vector plasmid pBin19 (Bevan, Nucl. Acids Research 12, (1984), 8711-8721) and was constructed by Höfgen and Willmitzer (Plant Sci. 66, (1990), 221-230). Starting from plasmid pB33-Kan, the *EcoRI-HindIII* fragment comprising the B33 promoter, a portion of the polylinker and the ocs terminator from pB33-Kan were cleaved and ligated into the vector pBIB-Hyg (Becker, Nucleic Acids Res. 18 (1), (1990), 203) which had been cleaved correspondingly. As a result, pB33-Hyg was obtained.

Then, an approximately 2000 bp-*Asp718* fragment of the plasmid pRL1 which contains the nucleotide sequence of about +2850 to about +4850 of the R1 cDNA from *Solanum tuberosum* (Lorberth, Charakterisierung von RL1: ein neues Enzym des Stärkemetabolismus. Dissertation Freie Universität Berlin) in antisense orientation into the *Asp718* cleavage site of the plasmid described before. The

resulting plasmid was called pB33-aR1-Hyg. For constructing the plasmid pB33-aBE1-Kan, analogously to the aforementioned strategy, the promoter region of the patatin-class-I gene B33 from *Solanum tuberosum* – a *Sma*I/*Hind*III fragment which has a length of about 3100 bp and contains a partial cDNA for the BE1 enzyme from potato (Kossmann, Klonierung und funktionelle Analyse von Genen codierend für am Kohlenhydratstoffwechsel der Kartoffel beteiligte Proteine, Dissertation Technische Universität Berlin, (1992)) – was first blunted and inserted into the *Sma*I cleavage site of the vector pBinAR-Hyg (cf. above) in antisense orientation with regard to the B33 promoter.

After the transformation, different lines of transgenic potato plants could be identified by means of Western blot analysis, the tubers of said potatoes having a content of the R1 protein which was reduced significantly. Further analyses showed that isolated starch of the line 36 had the highest amylose content of all transformants examined independently of each other.

Plants of said line were then transformed with the plasmid pGSV71-aBE2-basta as described by Rocha-Sosa et al. (EMBO J. 8 (1989), 23-29).

Plasmid pGSV71-aBE2-basta was constructed by screening according to standard procedures a tuber-specific potato cDNA library with a DNA fragment which had been amplified by RT-PCR (primer 1 (SEQ ID No. 1): 5'-gggggtgttggttgacta and primer 2 (SEQ ID No. 2) 5'-cccttctcctcctaatacca; Stratagene ProSTAR™ HF Single-Tube RT-PCR system) with total-RNA from tubers as template. In this way, a DNA fragment which has a size of about 1250 bp (cf. SEQ ID No. 3) and which was then subcloned as a *EcoRV-Sma*I fragment into the *EcoRV* cleavage site of the cloning vector pSK-Pac (cf. above) and finally ligated as *Pac*I fragment into the expression vector ME5/6 (Figure 1) in antisense orientation. As a result, the plasmid pGSV71-aBE2-basta was obtained.

From plants which were obtained by the transformation with the plasmid pGSV71-aBE2-basta and which showed a reduced R1, BEI and BEII gene expression, said plants being called 203MH plants, tissue samples of tubers of the independent transformants were taken and their amylose content was determined (cf. methods). The starches of the independent lines the tubers of which had the highest amylose



content were used for further analysing the starch properties (cf. Example 2).

### Example 2

#### Analysis of the starch of plants having a reduced R1, BEI and BEII gene expression

The starch of different independent lines of the transformation 203 MH described in Example 1 was isolated from potato tubers. Then, the physico-chemical properties of this starch were analysed. The results of the characterisation of the modified starches are shown in Table 1 (Tab. 1) for an exemplary selection of certain plant lines.

no.	genotype	phos- phate in C6 (%)	amylose (%)	RVA max (%)	RVA min (%)	RVA fin (%)	RVA set (%)	RVA T (%)	gel strength, 60% (w/v) CaCl <sub>2</sub> (%)
1	Desiree (wild type)	100	22	100	100	100	100	100	100
3	203MH010	25	92	no pasting in H <sub>2</sub> O					approx. 17,000
5	203MH055	33	80						not measured
6	203MH080	31	91						approx. 10,600

**Tab. 1**

Legend:

R1= R1 enzyme, BEI= branching enzyme I, BEII= branching enzyme II, as= antisense

RVA= Rapid Visco Analyser, max= maximum viscosity, min= minimum viscosity

fin= viscosity at the end of the measurement, set= set back= difference of min und fin

T= pasting temperature

The values in % are related to the wild type (= 100%) except for the amylose content.

The distribution of the side chains of the amylopectin was analysed as described above. The following Table (Tab. 2) contains an overview of the proportions of the individual peak heights of the HPAEC chromatograms within the overall peak height of wild type plants (Desiree), of 072VL036 plants (potato plants having a reduced gene expression of the R1 and BEI gene) and of selected lines of the transformations 203MH (cf. Example 1: potato plants having a reduced gene expression of the R1, BEI and BEII gene):

no. of glucose units	proportion of the individual peak heights within the overall peak height of every potato line in %				
	Desi Mix	072VL036	203MH10	203MH61	203MH80
dp6	2.48	1.2	2.6	1.7	2.7
dp7	1.90	1.2	1.6	1.1	1.5
dp8	1.46	1.4	1.0	0.8	1.2
dp9	2.48	2.2	1.3	1.1	1.5
dp10	4.38	3.5	2.1	1.9	2.1
dp11	6.28	4.8	3.1	2.8	3.0
dp12	7.30	5.6	3.6	3.6	3.3
dp13	7.88	6.1	3.6	3.9	3.3
dp14	7.88	6.3	3.9	4.4	3.6
dp15	7.30	6.3	4.2	4.4	3.3
dp16	6.72	6.1	4.2	4.7	3.3
dp17	5.84	5.9	4.2	4.7	3.3
dp18	5.26	5.7	4.4	4.7	3.3
dp19	4.82	5.5	4.4	4.7	3.3
dp20	4.38	5.1	4.4	4.7	3.3
dp21	3.94	4.7	4.4	4.7	3.6
dp22	3.50	4.2	4.2	4.7	3.6
dp23	3.07	3.8	4.2	4.4	3.9
dp24	2.48	3.4	4.4	4.4	4.2
dp25	2.19	3.1	4.2	4.4	4.2
dp26	1.90	2.7	4.2	4.2	4.5

dp27	1.75	2.4	3.9	4.2	4.5
dp28	1.31	2.0	3.6	3.6	4.5
dp29	1.02	1.6	3.4	3.3	4.2
dp30	0.88	1.3	2.9	2.8	3.9
dp31	0.58	1.1	2.6	2.5	3.3
dp32	0.44	0.8	2.1	1.9	2.7
dp33	0.29	0.7	1.8	1.7	2.4
dp34	0.29	0.5	1.8	1.4	2.4
dp35	0.00	0.4	1.6	1.1	1.8
dp36	0.00	0.3	1.0	0.8	1.5
dp37	0.00	0.2	0.8	0.6	0.9
dp38	0.00	0.2	0.5	0.0	0.9
dp39	0.00	0.0	0.0	0.0	0.0
dp40	0.00	0.0	0.0	0.0	0.0
Total	100.00	100.0	100.0	100.0	100.0

Tab. 2

If the proportion of peak heights of the individual chain lengths (indicated in DP) in the overall peak height is compared, a considerable shift towards side chains having a DP>26 can be seen with regard to the distribution of the side chains of the amylopectin of the 203MH plants compared to the amylopectin of wild type plants and also to 072VL plants. If the mean is calculated from the relative proportions of the side chains having a DP of 26 to DP 31, the following values are obtained (Tab. 3):

	Desi Mix	072VL036	203MH10	203MH61	203MH80
mean of the relative proportions of DP 26 to DP 31	1.24	1.85	3.43	3.43	4.15
change in % compared to Desi Mix (=100%)	100	149	276	276	335

Tab. 3

The amylopectin of 203MH plants is characterised by an increased proportion of side

chains having a DP of 26 to DP 31 compared to amylopectin of wild type plants and also of 072VL plants.

Furthermore, the morphology of the starch granules was examined:

The surface of the starch granules of wild type plants appears smooth under the light microscope. The form of the granules is round to oval, no "internal structures" being noticeable. Moreover, a uniform distribution of the different granule sizes can be seen (cf. Fig. 2).

Starch granules of 072VL036 plants (Fig. 3) have a very heterogeneous appearance. Only some granules appear smooth, others have grooves, some show "bunch-of-grapes-like agglomerations". Other granules have cross-recess-like structures. The spectrum of granule sizes is broad, smaller granules making up a greater proportion than is the case with starch granules of wild type plants.

The morphology of the granules of the line 203MH010 (Fig. 4), too, is heterogeneous, though less apparent than in 072VL036 plants. The surface of almost all granules has grooves, most of the granules show bunch-of-grapes-like agglomerations. Sometimes, particles can be seen which look like fragments of these agglomerated structures. The size distribution is relatively broad, smaller granules dominate though.

Furthermore, the granule size was determined using a photo sedimentometer of the type "Lumosed" by Retsch GmbH, Germany.

The average granule size of both untreated starch samples and samples which, prior to the granule size determination, were subjected to an overall 3-minute mechanical fragmentation was measured (for conduction see above) (Tab. 4).

In addition, the proportion of starch granules having a size of  $<20\text{ }\mu\text{m}$  was determined (Tab. 5)

average granule size [ $\mu\text{m}$ ]

sample	untreated	mechan. fragmentation
Wt	23.86	22.27
072VL036	16.88	16.78
203MH010	14.78	11.31
203MH066	14.59	11.82
203MH080	14.31	12.14

Tab. 4

Proportion of granules < 20 $\mu\text{m}$  [%]

sample	untreated	mechan. fragmentation
Wt	51.7	49.3
072VL036	69.7	69.9
203MH010	85.8	92.9
203MH066	83.7	90.6
203MH080	88.4	91.1

Tab. 5

The results show that both the average granule size and the proportion in percent of starch granules < 20  $\mu\text{m}$  of the starches of the invention differ significantly from wild type starches as well as from starches derived from 072VL036 plants.

After mechanical fragmentation of the starches, these differences are even more significant than without mechanical treatment.

### Claims

1. A transgenic plant cell which has been genetically modified, wherein the genetic modification leads to the reduction of the activity of one or more R1 proteins occurring endogenously in the plant cell and to the reduction of the activity of one or more BEI proteins occurring endogenously in the plant cell and to the reduction of the activity of one or more BEII proteins occurring endogenously in the plant cell in comparison with corresponding plant cells of wild type plants, said cells not being genetically modified.
2. The transgenic plant cell according to claim 1, wherein the genetic modification consists of the introduction of one or more foreign nucleic acid molecules the presence and/or expression of which leads to the reduction of the activity of R1 and BEI and BEII proteins in comparison with corresponding plant cells of wild type plants, said cells not being genetically modified.
3. The transgenic plant cells according to claim 2, wherein said foreign nucleic acid molecules are selected from the group consisting of
  - a) DNA molecules which encode at least one antisense RNA leading to a reduction of the expression of endogenous genes encoding R1 proteins and/or BEI proteins and/or BEII proteins;
  - b) DNA molecules which, through a co-suppression effect, lead to the reduction of the expression of endogenous genes encoding R1 proteins and/or BEI proteins and/or BEII proteins;
  - c) DNA molecules encoding at least one ribozyme which specifically cleaves transcripts of endogenous genes encoding R1 proteins and/or BEI proteins and/or BEII proteins; and
  - d) nucleic acid molecules which are introduced by means of *in-vivo* mutagenesis and which lead to a mutation or an insertion of a heterologous sequence in the genes encoding endogenous R1 proteins and/or BEI proteins and/or BEII proteins, wherein the mutation or insertion leads to a reduction of the expression of genes encoding R1 proteins

- and/or BEI proteins and/or BEII proteins, or to the synthesis of inactive R1 and/or BEI and/or BEII proteins; and
- e) DNA molecules which simultaneously encode at least one antisense RNA and at least one sense RNA, wherein said antisense RNA and said sense RNA form a double-stranded RNA molecule which leads to a reduction of the expression of endogenous genes encoding R1 proteins and/or BEI proteins and/or BEII proteins.
4. The transgenic plant cell according to any one of claims 1 to 3 which synthesises a modified starch.
  5. The transgenic plant cell according to claim 4, wherein the modified starch has an amylose content of at least 75% and a reduced phosphate content in comparison with starch of corresponding plant cells of wild type plants, said cells not being genetically modified.
  6. The transgenic plant cell according to claim 4 or 5, wherein the modified starch is characterised in that it has a modified distribution of the side chains.
  7. A transgenic plant containing plant cells according to any one of claims 1 to 6.
  8. A method for producing a transgenic plant cell which synthesises a modified starch, wherein a plant cell is genetically modified by the introduction of one or more foreign nucleic acid molecules the presence and/or expression of which lead to the reduction of the activity of R1, BEI and BEII proteins in comparison with corresponding plant cells of wild type plants, said cells not being genetically modified.
  9. The method of claim 8, wherein the modified starch is characterised in that it has an amylose content of at least 75% and a reduced phosphate content in comparison with starch of corresponding wild type plants which are not genetically modified.

10. A method for producing a transgenic plant which synthesises a modified starch, wherein
  - a) a plant cell is genetically modified by introducing one or more foreign nucleic acid molecules the presence and/or expression of which lead to the reduction of the activity of R1, BEI and BEII proteins compared to corresponding plant cells of wild type plants, said cells not being genetically modified;
  - b) a plant is regenerated from the cell produced according to a); and
  - c) optionally further plants are produced from the plant produced according to step b).
11. The method according to claim 10, wherein the modified starch is characterised in that it has an amylose content of at least 75% and a reduced phosphate content in comparison with starch of corresponding wild type plants which are not genetically modified.
12. The transgenic plant obtainable by a method according to claim 10 or 11.
13. The transgenic plant according to claim 7 or 12 which is a starch-storing plant.
14. The transgenic plant according to claim 7 or 12 which is a potato plant.
15. Propagation material of plants according to any one of claims 7 or 12 to 14 containing plant cells according to any one of claims 1 to 6.
16. Use of one or more foreign nucleic acid molecules encoding proteins having the enzymatic activity of R1, BEI and BEII proteins or fragments thereof for the production of plant cells according to any one of claims 1 to 6 or of plants according to any one of claims 7 or 12 to 14.
17. Use of one or more foreign nucleic acid molecules for the production of plants



according to any one of claims 7 or 12 to 14, wherein the foreign nucleic acid molecule is a molecule or the foreign nucleic acid molecules are several molecules selected from the group consisting of:

- a) DNA molecules which encode at least one antisense RNA leading to a reduction of the expression of endogenous genes encoding R1 proteins and/or BEI proteins and/or BEII proteins;
  - b) DNA molecules which, through a co-suppression effect, lead to the reduction of the expression of endogenous genes encoding R1 proteins and/or BEI proteins and/or BEII proteins;
  - c) DNA molecules encoding at least one ribozyme which specifically cleaves transcripts of endogenous genes encoding R1 proteins and/or BEI proteins and/or BEII proteins;
  - d) nucleic acid molecules which are introduced by means of *in-vivo* mutagenesis and which lead to a mutation or an insertion of a heterologous sequence in the genes encoding endogenous R1 proteins and/or BEI proteins and/or BEII proteins, wherein the mutation or insertion leads to a reduction of the expression of genes encoding R1 proteins and/or BEI proteins and/or BEII proteins, or the synthesis of inactive R1 and/or BEI and/or BEII proteins; and
  - e) DNA molecules which simultaneously encode at least one antisense RNA and at least one sense RNA, wherein said antisense RNA and said sense RNA form a double-stranded RNA molecule which leads to a reduction of the expression of endogenous genes encoding R1 proteins and/or BEI proteins and/or BEII proteins.
18. A composition containing at least one of the foreign nucleic acid molecules as defined in claim 17 which is suitable for the production of transgenic plant cells according to any one of claims 1 to 6 or transgenic plants according to any one of claims 7 or 12 to 14.
19. The composition according to claim 18, wherein the presence of said nucleic acid molecules in said plant cells leads to the reduction of the activity of R1 and

BEI and BEII proteins in comparison with corresponding plant cells of wild type plants, said cells not being genetically modified.

20. The composition according to claim 18 or 19, wherein the nucleic acid molecules are contained in a recombinant nucleic acid molecule.
21. A host cell containing a composition according to any one of claims 18 to 20.
22. A transgenic plant cell containing the composition according to any one of claims 18 to 20.
23. Starch obtainable from plant cells according to any one of claims 1 to 6 or 22 or from a plant according to any one of claims 7 or 12 to 14 or of propagation material according to claim 15.
24. Starch characterised in that it has an amylose content of at least 75% and a reduced phosphate content in comparison with starch from corresponding wild type plants which are not genetically modified.
25. Starch characterised in that it has an amylose content of at least 75% and a phosphate content in the C-6 position of the glucose monomers of up to 15 nmol C6-P mg<sup>-1</sup> starch.
26. Starch according to any one of claims 23 to 25 characterised in that it has a modified distribution of the side chains.
27. Starch according to any one of claims 23 to 26 which is a potato starch.
28. A method for producing a starch according to any one of claims 23 to 26 comprising the extraction of the starch from a plant according to any one of claims 7 or 12 to 14 and/or from starch-storing parts of such a plant and/or from a plant cell according to any one of claims 1 to 6 or 22 and/or propagation

material according to claim 15.

29. Starch obtainable by a method according to claim 28.





Figure 1



**Figure 2**




Figure 3




Figure 4

## 1.

## SEQUENCE LISTING

<110> Aventis CropScience GmbH  
 <120> Transgenic plants synthesising high amylose starch  
 <130> F 2027 PCT  
 <150> DE 10 12 8363.6  
 <151> 2001-06-12  
 <160> 10  
 <170> PatentIn version 3.1  
 <210> 1  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Oligonucleotide  
 <400> 1  
 ggggggtgttg gctttgacta 20  
 <210> 2  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Oligonucleotide  
 <400> 2  
 cccttctcct cctaattccca 20  
 <210> 3  
 <211> 1255  
 <212> DNA  
 <213> Solanum tuberosum  
 <220>  
 <221> CDS  
 <222> (2)..(928)  
 <223>  
 <400> 3  
 a ttt tgt att ccc gtt caa gat ggg ggt gtt ggc ttt gac tat cgg ctg 49  
 Phe Cys Ile Pro Val Gln Asp Gly Gly Val Gly Phe Asp Tyr Arg Leu  
 1 5 10 15  
 cat atg gca att gct gat aaa tgg att gag ttg ctc aag aaa cgg gat 97  
 His Met Ala Ile Ala Asp Lys Trp Ile Glu Leu Leu Lys Lys Arg Asp  
 20 25 30  
 gag gat tgg aga gtg ggt gat att gtt cat aca ctg aca aat aga aga 145



## 2

Glu	Asp	Trp	Arg	Val	Gly	Asp	Ile	Val	His	Thr	Leu	Thr	Asn	Arg	Arg	
	35						40					45				
tgg	tcg	gaa	aag	tgt	gtt	tca	tac	gct	gaa	agt	cat	gat	caa	gct	cta	193
Trp	Ser	Glu	Lys	Cys	Val	Ser	Tyr	Ala	Glu	Ser	His	Asp	Gln	Ala	Leu	
	50					55					60					
gtc	ggg	gat	aaa	act	ata	gca	ttc	tgg	ctg	atg	gac	aag	gat	atg	tat	241
Val	Gly	Asp	Lys	Thr	Ile	Ala	Phe	Trp	Leu	Met	Asp	Lys	Asp	Met	Tyr	
65					70					75					80	
gat	ttt	atg	gct	ttg	gat	aga	ccg	tca	aca	tca	tta	ata	gat	cgt	ggg	289
Asp	Phe	Met	Ala	Leu	Asp	Arg	Pro	Ser	Thr	Ser	Leu	Ile	Asp	Arg	Gly	
				85					90					95		
ata	gca	ttg	cac	aag	atg	att	agg	ctt	gta	act	atg	gga	tta	gga	gga	337
Ile	Ala	Leu	His	Lys	Met	Ile	Arg	Leu	Val	Thr	Met	Gly	Leu	Gly	Gly	
			100					105					110			
gaa	ggg	tac	cta	aat	ttc	atg	gga	aat	gaa	ttc	ggc	cac	cct	gag	tgg	385
Glu	Gly	Tyr	Leu	Asn	Phe	Met	Gly	Asn	Glu	Phe	Gly	His	Pro	Glu	Trp	
		115					120					125				
att	gat	ttc	cct	agg	gct	gaa	caa	cac	ctc	tct	gat	ggc	tca	gta	att	433
Ile	Asp	Phe	Pro	Arg	Ala	Glu	Gln	His	Leu	Ser	Asp	Gly	Ser	Val	Ile	
	130					135					140					
ccc	gga	aac	caa	ttc	agt	tat	gat	aaa	tgc	aga	cgg	aga	ttt	gac	ctg	481
Pro	Gly	Asn	Gln	Phe	Ser	Tyr	Asp	Lys	Cys	Arg	Arg	Arg	Phe	Asp	Leu	
145					150					155					160	
gga	gat	gca	gaa	tat	tta	aga	tac	cgt	ggg	ttg	caa	gaa	ttt	gac	cgg	529
Gly	Asp	Ala	Glu	Tyr	Leu	Arg	Tyr	Arg	Gly	Leu	Gln	Glu	Phe	Asp	Arg	
				165					170					175		
gct	atg	cag	tat	ctt	gaa	gat	aaa	tat	gag	ttt	atg	act	tca	gaa	cac	577
Ala	Met	Gln	Tyr	Leu	Glu	Asp	Lys	Tyr	Glu	Phe	Met	Thr	Ser	Glu	His	
			180					185					190			
cag	ttc	ata	tca	cga	aag	gat	gaa	gga	gat	agg	atg	att	gta	ttt	gaa	625
Gln	Phe	Ile	Ser	Arg	Lys	Asp	Glu	Gly	Asp	Arg	Met	Ile	Val	Phe	Glu	
		195					200					205				
aaa	gga	aac	cta	gtt	ttt	gtc	ttt	aat	ttt	cac	tgg	aca	aaa	agc	tat	673
Lys	Gly	Asn	Leu	Val	Phe	Val	Phe	Asn	Phe	His	Trp	Thr	Lys	Ser	Tyr	
	210					215					220					
tca	gac	tat	cgc	ata	ggc	tgc	ctg	aag	cct	gga	aaa	tac	aag	gtt	gcc	721
Ser	Asp	Tyr	Arg	Ile	Gly	Cys	Leu	Lys	Pro	Gly	Lys	Tyr	Lys	Val	Ala	
225					230					235					240	
ttg	gac	tca	gat	gat	cca	ctt	ttt	ggg	ggc	ttc	ggg	aga	att	gat	cat	769
Leu	Asp	Ser	Asp	Asp	Pro	Leu	Phe	Gly	Gly	Phe	Gly	Arg	Ile	Asp	His	
				245					250					255		
aat	gcc	gaa	tgt	ttc	acc	ttt	gaa	gga	tgg	tat	gat	gat	cgt	cct	cgt	817
Asn	Ala	Glu	Cys	Phe	Thr	Phe	Glu	Gly	Trp	Tyr	Asp	Asp	Arg	Pro	Arg	
			260					265					270			
tca	att	atg	gtg	tat	gca	cct	agt	aga	aca	gca	gtg	gtc	tat	gca	cta	865

## 3

Ser Ile Met Val Tyr Ala Pro Ser Arg Thr Ala Val Val Tyr Ala Leu  
 275 280 285

gta gac aaa gaa gaa gaa gaa gaa gaa gta gca gta gta gaa gaa gta 913  
 Val Asp Lys Glu Glu Glu Glu Glu Glu Val Ala Val Val Glu Glu Val  
 290 295 300

gta gta gaa gaa gaa tgaacgaact tgtgatcgcg ttgaaagatt tgaacgctac 968  
 Val Val Glu Glu Glu  
 305

atagagcttc ttgacgtatc tggcaatatt gcatcagtct tggcggaatt tcatgtgaca 1028

aaagggtttgc aattctttcc actattagta gtgcaacgat atacgcagag atgaagtgtc 1088

gaacaaacat atgtaaaatc gatgaattta tgtcgaatgc tgggacgggc ttcagcaggt 1148

tttgcttagt gagttctgta aattgtcatc tctttatatg tacagccaac tagaaatcaa 1208

ttatgtgaga cctaaaatac aataaccata aaatggaaat agtgctg 1255

<210> 4  
 <211> 309  
 <212> PRT  
 <213> Solanum tuberosum

<400> 4

Phe Cys Ile Pro Val Gln Asp Gly Gly Val Gly Phe Asp Tyr Arg Leu  
 1 5 10 15

His Met Ala Ile Ala Asp Lys Trp Ile Glu Leu Leu Lys Lys Arg Asp  
 20 25 30

Glu Asp Trp Arg Val Gly Asp Ile Val His Thr Leu Thr Asn Arg Arg  
 35 40 45

Trp Ser Glu Lys Cys Val Ser Tyr Ala Glu Ser His Asp Gln Ala Leu  
 50 55 60

Val Gly Asp Lys Thr Ile Ala Phe Trp Leu Met Asp Lys Asp Met Tyr  
 65 70 75 80

Asp Phe Met Ala Leu Asp Arg Pro Ser Thr Ser Leu Ile Asp Arg Gly  
 85 90 95

Ile Ala Leu His Lys Met Ile Arg Leu Val Thr Met Gly Leu Gly Gly  
 100 105 110

Glu Gly Tyr Leu Asn Phe Met Gly Asn Glu Phe Gly His Pro Glu Trp  
 115 120 125

## 4

Ile Asp Phe Pro Arg Ala Glu Gln His Leu Ser Asp Gly Ser Val Ile  
 130 135 140  
 Pro Gly Asn Gln Phe Ser Tyr Asp Lys Cys Arg Arg Arg Phe Asp Leu  
 145 150 155 160  
 Gly Asp Ala Glu Tyr Leu Arg Tyr Arg Gly Leu Gln Glu Phe Asp Arg  
 165 170 175  
 Ala Met Gln Tyr Leu Glu Asp Lys Tyr Glu Phe Met Thr Ser Glu His  
 180 185 190  
 Gln Phe Ile Ser Arg Lys Asp Glu Gly Asp Arg Met Ile Val Phe Glu  
 195 200 205  
 Lys Gly Asn Leu Val Phe Val Phe Asn Phe His Trp Thr Lys Ser Tyr  
 210 215 220  
 Ser Asp Tyr Arg Ile Gly Cys Leu Lys Pro Gly Lys Tyr Lys Val Ala  
 225 230 235 240  
 Leu Asp Ser Asp Asp Pro Leu Phe Gly Gly Phe Gly Arg Ile Asp His  
 245 250 255  
 Asn Ala Glu Cys Phe Thr Phe Glu Gly Trp Tyr Asp Asp Arg Pro Arg  
 260 265 270  
 Ser Ile Met Val Tyr Ala Pro Ser Arg Thr Ala Val Val Tyr Ala Leu  
 275 280 285  
 Val Asp Lys Glu Glu Glu Glu Glu Glu Val Ala Val Val Glu Glu Val  
 290 295 300  
 Val Val Glu Glu Glu  
 305

<210> 5  
 <211> 5061  
 <212> DNA  
 <213> Solanum tuberosum

<220>  
 <221> CDS  
 <222> (216)..(4607)  
 <223>

5,

<400> 5  
gaattgtaat acgactcact atagggcgaa ttgggtaccg ggccccccct cgagggtcgac 60  
ggtatcgata agcttgatat cgaattcgcg gccgcttttg ctctgtgaat tcattctcat 120  
cgaattttctc gacgcttctt cgctaatttc ctctgttactt cactagaaat cgacgtttct 180  
agctgaactt gagtgaatta agccagtggg aggat atg agt aat tcc tta ggg 233  
Met Ser Asn Ser Leu Gly  
1 5  
aat aac ttg ctg tac cag gga ttc cta acc tca aca gtg ttg gaa cat 281  
Asn Asn Leu Leu Tyr Gln Gly Phe Leu Thr Ser Thr Val Leu Glu His  
10 15 20  
aaa agt aga atc agt cct cct tgt gtt gga ggc aat tct ttg ttt caa 329  
Lys Ser Arg Ile Ser Pro Pro Cys Val Gly Gly Asn Ser Leu Phe Gln  
25 30 35  
caa caa gtg atc tcg aaa tca cct tta tca act gag ttt cga ggt aac 377  
Gln Gln Val Ile Ser Lys Ser Pro Leu Ser Thr Glu Phe Arg Gly Asn  
40 45 50  
agg tta aag gtg cag aaa aag aaa ata cct atg gga aag aac cgt gct 425  
Arg Leu Lys Val Gln Lys Lys Lys Ile Pro Met Gly Lys Asn Arg Ala  
55 60 65 70  
ttt tct agt tct cct cat gct gta ctt acc act gat acc tct tct gag 473  
Phe Ser Ser Ser Pro His Ala Val Leu Thr Thr Asp Thr Ser Ser Glu  
75 80 85  
cta gca gaa aag ttc agt cta gaa ggg aat att gag cta cag gtt gat 521  
Leu Ala Glu Lys Phe Ser Leu Glu Gly Asn Ile Glu Leu Gln Val Asp  
90 95 100  
gtt agg cct ccc act tca ggt gat gtg tcc ttt gtg gat ttt caa gct 569  
Val Arg Pro Pro Thr Ser Gly Asp Val Ser Phe Val Asp Phe Gln Ala  
105 110 115  
aca aat ggt agt gat aaa ctg ttt ttg cac tgg ggg gca gta aag ttc 617  
Thr Asn Gly Ser Asp Lys Leu Phe Leu His Trp Gly Ala Val Lys Phe  
120 125 130  
gga aaa gaa aca tgg tct ctt cct aat gat cgt cca gat ggg acc aaa 665  
Gly Lys Glu Thr Trp Ser Leu Pro Asn Asp Arg Pro Asp Gly Thr Lys  
135 140 145 150  
gtg tac aag aac aaa gca ctt aga act cca ttt gtt aaa tct ggc tct 713  
Val Tyr Lys Asn Lys Ala Leu Arg Thr Pro Phe Val Lys Ser Gly Ser  
155 160 165  
aac tcc atc ctg aga ctg gag ata cgg gac act gct atc gaa gct att 761  
Asn Ser Ile Leu Arg Leu Glu Ile Arg Asp Thr Ala Ile Glu Ala Ile  
170 175 180  
gag ttt ctg ata tac gat gaa gcc tac gat aaa tgg ata aag aat aat 809  
Glu Phe Leu Ile Tyr Asp Glu Ala Tyr Asp Lys Trp Ile Lys Asn Asn  
185 190 195  
ggg ggc aat ttt cgt gtc aaa ttg tca aga aaa gag ata cga ggc cca 857

Gly	Gly	Asn	Phe	Arg	Val	Lys	Leu	Ser	Arg	Lys	Glu	Ile	Arg	Gly	Pro	
200						205					210					
gat	ggt	tca	ggt	cct	gag	gag	ctt	gta	cag	atc	caa	tca	tat	ttg	agg	905
Asp	Val	Ser	Val	Pro	Glu	Glu	Leu	Val	Gln	Ile	Gln	Ser	Tyr	Leu	Arg	
215					220				225					230		
tgg	gag	agg	aag	gga	aaa	cag	aat	tac	acc	cct	gag	aaa	gag	aag	gag	953
Trp	Glu	Arg	Lys	Gly	Lys	Gln	Asn	Tyr	Thr	Pro	Glu	Lys	Glu	Lys	Glu	
			235					240					245			
gaa	tat	gag	gct	gct	cga	act	gag	cta	cag	gag	gaa	ata	gct	cgt	ggt	1001
Glu	Tyr	Glu	Ala	Ala	Arg	Thr	Glu	Leu	Gln	Glu	Glu	Ile	Ala	Arg	Gly	
			250					255					260			
gct	tcc	ata	cag	gac	att	cga	gca	agg	cta	aca	aaa	act	aat	gat	aaa	1049
Ala	Ser	Ile	Gln	Asp	Ile	Arg	Ala	Arg	Leu	Thr	Lys	Thr	Asn	Asp	Lys	
		265					270					275				
agt	caa	agc	aaa	gaa	gag	cct	ctt	cat	gta	aca	aag	agt	gaa	ata	cct	1097
Ser	Gln	Ser	Lys	Glu	Glu	Pro	Leu	His	Val	Thr	Lys	Ser	Glu	Ile	Pro	
	280					285					290					
gat	gac	ctt	gcc	caa	gca	caa	gct	tac	att	agg	tgg	gag	aaa	gca	gga	1145
Asp	Asp	Leu	Ala	Gln	Ala	Gln	Ala	Tyr	Ile	Arg	Trp	Glu	Lys	Ala	Gly	
295				300						305				310		
aag	ccg	aac	tat	cct	cca	gaa	aag	caa	att	gaa	gaa	ctc	gaa	gaa	gca	1193
Lys	Pro	Asn	Tyr	Pro	Pro	Glu	Lys	Gln	Ile	Glu	Glu	Leu	Glu	Glu	Ala	
			315					320					325			
aga	aga	gaa	ttg	caa	ctt	gag	ctt	gag	aaa	ggc	att	acc	ctt	gat	gag	1241
Arg	Arg	Glu	Leu	Gln	Leu	Glu	Leu	Glu	Lys	Gly	Ile	Thr	Leu	Asp	Glu	
		330						335					340			
ttg	cgg	aaa	aag	att	aca	aaa	ggg	gag	ata	aaa	act	aag	gcg	gaa	aag	1289
Leu	Arg	Lys	Lys	Ile	Thr	Lys	Gly	Glu	Ile	Lys	Thr	Lys	Ala	Glu	Lys	
		345					350					355				
cac	gtg	aaa	aga	agc	tct	ttt	gcc	gtt	gaa	aga	atc	caa	aga	aag	aag	1337
His	Val	Lys	Arg	Ser	Ser	Phe	Ala	Val	Glu	Arg	Ile	Gln	Arg	Lys	Lys	
	360					365					370					
aga	gac	ttt	ggg	cag	ctt	att	aat	aag	tat	cct	tcc	agt	cct	gca	gta	1385
Arg	Asp	Phe	Gly	Gln	Leu	Ile	Asn	Lys	Tyr	Pro	Ser	Ser	Pro	Ala	Val	
	375				380					385				390		
caa	gta	caa	aag	gtc	ttg	gaa	gaa	cca	cca	gcc	tta	tct	aaa	att	aag	1433
Gln	Val	Gln	Lys	Val	Leu	Glu	Glu	Pro	Pro	Ala	Leu	Ser	Lys	Ile	Lys	
			395					400					405			
ctg	tat	gcc	aag	gag	aag	gag	gag	cag	att	gat	gat	ccg	atc	ctt	aat	1481
Leu	Tyr	Ala	Lys	Glu	Lys	Glu	Glu	Gln	Ile	Asp	Asp	Pro	Ile	Leu	Asn	
		410						415					420			
aaa	aag	atc	ttt	aag	gtc	gat	gat	ggg	gag	cta	ctg	gta	ctg	gta	gca	1529
Lys	Lys	Ile	Phe	Lys	Val	Asp	Asp	Gly	Glu	Leu	Leu	Val	Leu	Val	Ala	
		425					430					435				
aag	tcc	tct	ggg	aag	aca	aaa	gta	cat	ata	gct	aca	gat	ctg	aat	cag	1577

7.

Lys	Ser	Ser	Gly	Lys	Thr	Lys	Val	His	Ile	Ala	Thr	Asp	Leu	Asn	Gln	
440						445					450					
cca	att	act	ctt	cac	tgg	gca	tta	tcc	aaa	agt	cgt	gga	gag	tgg	atg	1625
Pro	Ile	Thr	Leu	His	Trp	Ala	Leu	Ser	Lys	Ser	Arg	Gly	Glu	Trp	Met	
455					460					465					470	
gta	cca	cct	tca	agc	ata	ttg	cct	cct	gga	tca	att	att	tta	gac	aag	1673
Val	Pro	Pro	Ser	Ser	Ile	Leu	Pro	Pro	Gly	Ser	Ile	Ile	Leu	Asp	Lys	
				475					480					485		
gct	gcc	gaa	aca	cct	ttt	tcc	gcc	agt	tct	tct	gat	ggt	cta	act	tct	1721
Ala	Ala	Glu	Thr	Pro	Phe	Ser	Ala	Ser	Ser	Ser	Asp	Gly	Leu	Thr	Ser	
			490					495					500			
aag	gta	caa	tct	ttg	gat	ata	gta	att	gaa	gat	ggc	aat	ttt	gtg	ggg	1769
Lys	Val	Gln	Ser	Leu	Asp	Ile	Val	Ile	Glu	Asp	Gly	Asn	Phe	Val	Gly	
		505					510					515				
atg	cca	ttt	gtt	ctt	ttg	tct	ggt	gaa	aaa	tgg	att	aag	aac	caa	ggg	1817
Met	Pro	Phe	Val	Leu	Leu	Ser	Gly	Glu	Lys	Trp	Ile	Lys	Asn	Gln	Gly	
	520					525				530						
tcg	gat	ttc	tat	gtt	gac	ttc	agt	gct	gca	tcc	aaa	tta	gca	ctc	aag	1865
Ser	Asp	Phe	Tyr	Val	Asp	Phe	Ser	Ala	Ala	Ser	Lys	Leu	Ala	Leu	Lys	
535					540					545					550	
gct	gct	ggg	gat	ggc	agt	gga	act	gca	aag	tct	tta	ctg	gat	aaa	ata	1913
Ala	Ala	Gly	Asp	Gly	Ser	Gly	Thr	Ala	Lys	Ser	Leu	Leu	Asp	Lys	Ile	
				555					560					565		
gca	gat	atg	gaa	agt	gag	gct	cag	aag	tca	ttt	atg	cac	cgg	ttt	aat	1961
Ala	Asp	Met	Glu	Ser	Glu	Ala	Gln	Lys	Ser	Phe	Met	His	Arg	Phe	Asn	
			570					575					580			
att	gct	gct	gac	ttg	ata	gaa	gat	gcc	act	agt	gct	ggt	gaa	ctt	ggt	2009
Ile	Ala	Ala	Asp	Leu	Ile	Glu	Asp	Ala	Thr	Ser	Ala	Gly	Glu	Leu	Gly	
			585				590					595				
ttt	act	gga	att	ctt	gta	tgg	atg	agg	ttc	atg	gct	aca	agg	caa	ctg	2057
Phe	Thr	Gly	Ile	Leu	Val	Trp	Met	Arg	Phe	Met	Ala	Thr	Arg	Gln	Leu	
	600					605				610						
ata	tgg	aac	aaa	aac	tat	aac	gta	aaa	cca	cgt	gaa	ata	agc	aag	gct	2105
Ile	Trp	Asn	Lys	Asn	Tyr	Asn	Val	Lys	Pro	Arg	Glu	Ile	Ser	Lys	Ala	
615					620					625					630	
cag	gac	aga	ctt	aca	gac	ttg	ttg	cag	aat	gct	ttc	acc	agt	cac	cct	2153
Gln	Asp	Arg	Leu	Thr	Asp	Leu	Leu	Gln	Asn	Ala	Phe	Thr	Ser	His	Pro	
				635					640					645		
caa	tac	cgt	gaa	att	ttg	cgg	atg	att	atg	tca	act	gtt	gga	cgt	gga	2201
Gln	Tyr	Arg	Glu	Ile	Leu	Arg	Met	Ile	Met	Ser	Thr	Val	Gly	Arg	Gly	
			650					655					660			
ggt	gaa	ggg	gat	gta	gga	cag	cga	att	agg	gat	gaa	att	ttg	gtc	atc	2249
Gly	Glu	Gly	Asp	Val	Gly	Gln	Arg	Ile	Arg	Asp	Glu	Ile	Leu	Val	Ile	
		665					670					675				
cag	agg	aaa	aat	gac	tgc	aag	ggt	ggt	atg	atg	gaa	gaa	tgg	cat	cag	2297

## 8.

Gln	Arg	Lys	Asn	Asp	Cys	Lys	Gly	Gly	Met	Met	Glu	Glu	Trp	His	Gln	
680						685					690					
aaa	ttg	cat	aat	aat	act	agt	cct	gat	gat	gtt	gtg	atc	tgt	cag	gca	2345
Lys	Leu	His	Asn	Asn	Thr	Ser	Pro	Asp	Asp	Val	Val	Ile	Cys	Gln	Ala	
695					700					705					710	
ttg	att	gac	tac	atc	aag	agt	gat	ttt	gat	ctt	ggt	gtt	tat	tgg	aaa	2393
Leu	Ile	Asp	Tyr	Ile	Lys	Ser	Asp	Phe	Asp	Leu	Gly	Val	Tyr	Trp	Lys	
				715					720					725		
acc	ctg	aat	gag	aac	gga	ata	aca	aaa	gag	cgt	ctt	ttg	agt	tat	gac	2441
Thr	Leu	Asn	Glu	Asn	Gly	Ile	Thr	Lys	Glu	Arg	Leu	Leu	Ser	Tyr	Asp	
			730					735					740			
cgt	gct	atc	cat	tct	gaa	ccg	aat	ttt	aga	gga	gat	caa	aag	aat	ggt	2489
Arg	Ala	Ile	His	Ser	Glu	Pro	Asn	Phe	Arg	Gly	Asp	Gln	Lys	Asn	Gly	
		745					750					755				
ctt	ttg	cgt	gat	tta	ggt	cac	tat	atg	aga	aca	ttg	aag	gct	gtt	cat	2537
Leu	Leu	Arg	Asp	Leu	Gly	His	Tyr	Met	Arg	Thr	Leu	Lys	Ala	Val	His	
	760					765					770					
tca	ggt	gca	gat	ctt	gag	tct	gct	att	gca	aac	tgc	atg	ggc	tac	aaa	2585
Ser	Gly	Ala	Asp	Leu	Glu	Ser	Ala	Ile	Ala	Asn	Cys	Met	Gly	Tyr	Lys	
775					780				785						790	
act	gag	gga	gaa	ggc	ttt	atg	gtt	gga	gtc	cag	ata	aat	cct	gta	tca	2633
Thr	Glu	Gly	Glu	Gly	Phe	Met	Val	Gly	Val	Gln	Ile	Asn	Pro	Val	Ser	
				795					800					805		
ggc	ttg	cca	tct	ggc	ttt	cag	ggc	ctc	ctc	cat	ttt	gtc	tta	gac	cat	2681
Gly	Leu	Pro	Ser	Gly	Phe	Gln	Gly	Leu	Leu	His	Phe	Val	Leu	Asp	His	
			810					815					820			
gtg	gaa	gat	aaa	aat	gtg	gaa	act	ctt	ctt	gag	gga	ttg	cta	gag	gct	2729
Val	Glu	Asp	Lys	Asn	Val	Glu	Thr	Leu	Leu	Glu	Gly	Leu	Leu	Glu	Ala	
		825					830					835				
cgt	gag	gag	ctt	agg	ccc	ttg	ctt	ctc	aaa	cca	aac	aac	cgt	cta	aag	2777
Arg	Glu	Glu	Leu	Arg	Pro	Leu	Leu	Leu	Lys	Pro	Asn	Asn	Arg	Leu	Lys	
	840					845					850					
gat	ctg	ctg	ttt	ttg	gac	ata	gca	ctt	gat	tct	aca	gtt	aga	aca	gca	2825
Asp	Leu	Leu	Phe	Leu	Asp	Ile	Ala	Leu	Asp	Ser	Thr	Val	Arg	Thr	Ala	
855					860					865					870	
gta	gaa	agg	gga	tat	gaa	gaa	ttg	aac	aac	gct	aat	cct	gag	aaa	atc	2873
Val	Glu	Arg	Gly	Tyr	Glu	Glu	Leu	Asn	Asn	Ala	Asn	Pro	Glu	Lys	Ile	
				875					880					885		
atg	tac	ttc	atc	tcc	ctc	gtt	ctt	gaa	aat	ctc	gca	ctc	tct	gtg	gac	2921
Met	Tyr	Phe	Ile	Ser	Leu	Val	Leu	Glu	Asn	Leu	Ala	Leu	Ser	Val	Asp	
			890					895					900			
gat	aat	gaa	gat	ctt	gtt	tat	tgc	ttg	aag	gga	tgg	aat	caa	gct	ctt	2969
Asp	Asn	Glu	Asp	Leu	Val	Tyr	Cys	Leu	Lys	Gly	Trp	Asn	Gln	Ala	Leu	
		905					910					915				
tca	atg	tcc	aat	ggt	gga	gac	aac	cat	tgg	gct	tta	ttt	gca	aaa	gct	3017

## 9

Ser	Met	Ser	Asn	Gly	Gly	Asp	Asn	His	Trp	Ala	Leu	Phe	Ala	Lys	Ala		
920						925					930						
gta	ctt	gac	aga	atc	cgt	ctt	gca	ctt	gca	agc	aag	gca	gag	tgg	tac	3065	
Val	Leu	Asp	Arg	Ile	Arg	Leu	Ala	Leu	Ala	Ser	Lys	Ala	Glu	Trp	Tyr		
935					940					945					950		
cat	cac	tta	ttg	cag	cca	tct	gcc	gaa	tat	cta	gga	tca	atc	ctt	ggg	3113	
His	His	Leu	Leu	Gln	Pro	Ser	Ala	Glu	Tyr	Leu	Gly	Ser	Ile	Leu	Gly		
				955					960						965		
gtg	gac	caa	ttg	gct	ttg	aac	ata	ttt	act	gaa	gaa	att	ata	cgt	gct	3161	
Val	Asp	Gln	Trp	Ala	Leu	Asn	Ile	Phe	Thr	Glu	Glu	Ile	Ile	Arg	Ala		
			970					975							980		
gga	tca	gca	gct	tca	tta	tcc	tct	ctt	ctt	aat	aga	ctc	gat	ccc	gtg	3209	
Gly	Ser	Ala	Ala	Ser	Leu	Ser	Ser	Leu	Leu	Asn	Arg	Leu	Asp	Pro	Val		
		985					990					995					
ctt	cgg	aaa	act	gca	aat	cta	gga	agt	tgg	cag	att	atc	agt	cca		3254	
Leu	Arg	Lys	Thr	Ala	Asn	Leu	Gly	Ser	Trp	Gln	Ile	Ile	Ser	Pro			
	1000					1005						1010					
gtt	gaa	gcc	gtt	gga	tat	gtt	gtc	gtt	gtg	gat	gag	ttg	ctt	tca		3299	
Val	Glu	Ala	Val	Gly	Tyr	Val	Val	Val	Val	Asp	Glu	Leu	Leu	Ser			
	1015					1020					1025						
gtt	cag	aat	gaa	atc	tac	aag	aag	ccc	acg	atc	tta	gta	gca	aac		3344	
Val	Gln	Asn	Glu	Ile	Tyr	Lys	Lys	Pro	Thr	Ile	Leu	Val	Ala	Asn			
	1030					1035					1040						
tct	gtt	aaa	gga	gag	gag	gaa	att	cct	gat	ggt	gct	gtt	gcc	ctg		3389	
Ser	Val	Lys	Gly	Glu	Glu	Glu	Ile	Pro	Asp	Gly	Ala	Val	Ala	Leu			
	1045					1050					1055						
ata	aca	cca	gac	atg	cca	gat	gtt	ctt	tca	cat	gtt	tct	gtt	cga		3434	
Ile	Thr	Pro	Asp	Met	Pro	Asp	Val	Leu	Ser	His	Val	Ser	Val	Arg			
	1060					1065					1070						
gct	aga	aat	ggg	aag	gtt	tgc	ttt	gct	aca	tgc	ttt	gat	ccc	aat		3479	
Ala	Arg	Asn	Gly	Lys	Val	Cys	Phe	Ala	Thr	Cys	Phe	Asp	Pro	Asn			
	1075					1080					1085						
ata	ttg	gct	gac	ctc	caa	gca	aag	gaa	gga	agg	att	ttg	ctc	tta		3524	
Ile	Leu	Ala	Asp	Leu	Gln	Ala	Lys	Glu	Gly	Arg	Ile	Leu	Leu	Leu			
	1090					1095					1100						
aag	cct	aca	cct	tca	gac	ata	atc	tat	agt	gag	gtg	aat	gag	att		3569	
Lys	Pro	Thr	Pro	Ser	Asp	Ile	Ile	Tyr	Ser	Glu	Val	Asn	Glu	Ile			
	1105					1110					1115						
gag	ctc	caa	agt	tca	agt	aac	ttg	gta	gaa	gct	gaa	act	tca	gca		3614	
Glu	Leu	Gln	Ser	Ser	Ser	Asn	Leu	Val	Glu	Ala	Glu	Thr	Ser	Ala			
	1120					1125					1130						
aca	ctt	aga	ttg	gtg	aaa	aag	caa	ttt	ggt	ggt	tgt	tac	gca	ata		3659	
Thr	Leu	Arg	Leu	Val	Lys	Lys	Gln	Phe	Gly	Gly	Cys	Tyr	Ala	Ile			
	1135					1140					1145						
tca	gca	gat	gaa	ttc	aca	agt	gaa	atg	gtt	gga	gct	aaa	tca	cgt		3704	



## 10.

Ser	Ala	Asp	Glu	Phe	Thr	Ser	Glu	Met	Val	Gly	Ala	Lys	Ser	Arg	
1150						1155					1160				
aat	att	gca	tat	ctg	aaa	gga	aaa	gtg	cct	tcc	tcg	gtg	gga	att	3749
Asn	Ile	Ala	Tyr	Leu	Lys	Gly	Lys	Val	Pro	Ser	Ser	Val	Gly	Ile	
1165						1170					1175				
cct	acg	tca	gta	gct	ctt	cca	ttt	gga	gtc	ttt	gag	aaa	gta	ctt	3794
Pro	Thr	Ser	Val	Ala	Leu	Pro	Phe	Gly	Val	Phe	Glu	Lys	Val	Leu	
1180						1185					1190				
tca	gac	gac	ata	aat	cag	gga	gtg	gca	aaa	gag	ttg	caa	att	ctg	3839
Ser	Asp	Asp	Ile	Asn	Gln	Gly	Val	Ala	Lys	Glu	Leu	Gln	Ile	Leu	
1195						1200					1205				
acg	aaa	aaa	cta	tct	gaa	gga	gac	ttc	agc	gct	ctt	ggg	gaa	att	3884
Thr	Lys	Lys	Leu	Ser	Glu	Gly	Asp	Phe	Ser	Ala	Leu	Gly	Glu	Ile	
1210						1215					1220				
cgc	aca	acg	att	tta	gat	ctt	tca	gca	cca	gct	caa	ttg	gtc	aaa	3929
Arg	Thr	Thr	Ile	Leu	Asp	Leu	Ser	Ala	Pro	Ala	Gln	Leu	Val	Lys	
1225						1230					1235				
gag	ctg	aag	gaa	aag	atg	cag	ggg	tct	ggc	atg	cct	tgg	cct	ggg	3974
Glu	Leu	Lys	Glu	Lys	Met	Gln	Gly	Ser	Gly	Met	Pro	Trp	Pro	Gly	
1240						1245					1250				
gat	gaa	ggg	cca	aag	cgg	tgg	gaa	caa	gca	tgg	atg	gcc	ata	aaa	4019
Asp	Glu	Gly	Pro	Lys	Arg	Trp	Glu	Gln	Ala	Trp	Met	Ala	Ile	Lys	
1255						1260					1265				
aag	gtg	tgg	gct	tca	aaa	tgg	aat	gag	aga	gca	tac	ttc	agc	aca	4064
Lys	Val	Trp	Ala	Ser	Lys	Trp	Asn	Glu	Arg	Ala	Tyr	Phe	Ser	Thr	
1270						1275					1280				
agg	aag	gtg	aaa	ctg	gat	cat	gac	tat	ctg	tgc	atg	gct	gtc	ctt	4109
Arg	Lys	Val	Lys	Leu	Asp	His	Asp	Tyr	Leu	Cys	Met	Ala	Val	Leu	
1285						1290					1295				
gtt	caa	gaa	ata	ata	aat	gct	gat	tat	gca	ttt	gtc	att	cac	aca	4154
Val	Gln	Glu	Ile	Ile	Asn	Ala	Asp	Tyr	Ala	Phe	Val	Ile	His	Thr	
1300						1305					1310				
acc	aac	cca	tct	tcc	gga	gac	gac	tca	gaa	ata	tat	gcc	gag	gtg	4199
Thr	Asn	Pro	Ser	Ser	Gly	Asp	Asp	Ser	Glu	Ile	Tyr	Ala	Glu	Val	
1315						1320					1325				
gtc	agg	ggc	ctt	ggg	gaa	aca	ctt	gtt	gga	gct	tac	cca	gga	cgt	4244
Val	Arg	Gly	Leu	Gly	Glu	Thr	Leu	Val	Gly	Ala	Tyr	Pro	Gly	Arg	
1330						1335					1340				
gct	ttg	agt	ttt	atc	tgc	aag	aaa	aag	gat	ctc	aac	tct	cct	caa	4289
Ala	Leu	Ser	Phe	Ile	Cys	Lys	Lys	Lys	Asp	Leu	Asn	Ser	Pro	Gln	
1345						1350					1355				
gtg	tta	ggg	tac	cca	agc	aaa	ccg	atc	ggc	ctt	ttc	ata	aaa	aga	4334
Val	Leu	Gly	Tyr	Pro	Ser	Lys	Pro	Ile	Gly	Leu	Phe	Ile	Lys	Arg	
1360						1365					1370				
tct	atc	atc	ttc	cga	tct	gat	tcc	aat	ggg	gaa	gat	ttg	gaa	ggg	4379

Ser Ile Ile Phe Arg Ser Asp Ser Asn Gly Glu Asp Leu Glu Gly  
 1375 1380 1385  
 tat gcc ggt gct ggc ctc tac gac agt gta cca atg gat gag gag 4424  
 Tyr Ala Gly Ala Gly Leu Tyr Asp Ser Val Pro Met Asp Glu Glu  
 1390 1395 1400  
 gaa aaa gtt gta att gat tac tct tcc gac cca ttg ata act gat 4469  
 Glu Lys Val Val Ile Asp Tyr Ser Ser Asp Pro Leu Ile Thr Asp  
 1405 1410 1415  
 ggt aac ttc cgc cag aca atc ctg tcc aac att gct cgt gct gga 4514  
 Gly Asn Phe Arg Gln Thr Ile Leu Ser Asn Ile Ala Arg Ala Gly  
 1420 1425 1430  
 cat gct atc gag gag cta tat ggc tct cct caa gac atc gag ggt 4559  
 His Ala Ile Glu Glu Leu Tyr Gly Ser Pro Gln Asp Ile Glu Gly  
 1435 1440 1445  
 gta gtg agg gat gga aag att tat gtc gtt cag aca aga cct cag 4604  
 Val Val Arg Asp Gly Lys Ile Tyr Val Val Gln Thr Arg Pro Gln  
 1450 1455 1460  
 atg tgatcatatt ctcgttgtat gttgttcaga gaagaccata gatgtgatca 4657  
 Met  
 tattctcatg gtatcagatc tgtgaccact tacctcccat gaagttgcct gtatgattat 4717  
 acgtgatcca aagccatcac atcatgttca ccttcagcta ttggaggaga agtgagaagt 4777  
 aggaattgca atatgaggaa taataagaaa aactttgtag aagttaaatt agctgggtat 4837  
 gatataggga gaaatgtgta aacattgtac tatatatagt atacacacgc attatgtatt 4897  
 tgcattatgc actgaataat atcgagcat caaagaagaa atcctttgag tggtttcaat 4957  
 tgccgcggcc gcgaattcct gcagcccggtt ggatccacta gttctagagc ggccgccacc 5017  
 gcggtggagc tccagctttt gttcccttta gtgagggtta attt 5061  
 <210> 6  
 <211> 1464  
 <212> PRT  
 <213> Solanum tuberosum  
 <400> 6  
 Met Ser Asn Ser Leu Gly Asn Asn Leu Leu Tyr Gln Gly Phe Leu Thr  
 1 5 10 15  
 Ser Thr Val Leu Glu His Lys Ser Arg Ile Ser Pro Pro Cys Val Gly  
 20 25 30  
 Gly Asn Ser Leu Phe Gln Gln Gln Val Ile Ser Lys Ser Pro Leu Ser  
 35 40 45

## 12.

Thr Glu Phe Arg Gly Asn Arg Leu Lys Val Gln Lys Lys Lys Ile Pro  
 50 55 60

Met Gly Lys Asn Arg Ala Phe Ser Ser Ser Pro His Ala Val Leu Thr  
 65 70 75 80

Thr Asp Thr Ser Ser Glu Leu Ala Glu Lys Phe Ser Leu Glu Gly Asn  
 85 90 95

Ile Glu Leu Gln Val Asp Val Arg Pro Pro Thr Ser Gly Asp Val Ser  
 100 105 110

Phe Val Asp Phe Gln Ala Thr Asn Gly Ser Asp Lys Leu Phe Leu His  
 115 120 125

Trp Gly Ala Val Lys Phe Gly Lys Glu Thr Trp Ser Leu Pro Asn Asp  
 130 135 140

Arg Pro Asp Gly Thr Lys Val Tyr Lys Asn Lys Ala Leu Arg Thr Pro  
 145 150 155 160

Phe Val Lys Ser Gly Ser Asn Ser Ile Leu Arg Leu Glu Ile Arg Asp  
 165 170 175

Thr Ala Ile Glu Ala Ile Glu Phe Leu Ile Tyr Asp Glu Ala Tyr Asp  
 180 185 190

Lys Trp Ile Lys Asn Asn Gly Gly Asn Phe Arg Val Lys Leu Ser Arg  
 195 200 205

Lys Glu Ile Arg Gly Pro Asp Val Ser Val Pro Glu Glu Leu Val Gln  
 210 215 220

Ile Gln Ser Tyr Leu Arg Trp Glu Arg Lys Gly Lys Gln Asn Tyr Thr  
 225 230 235 240

Pro Glu Lys Glu Lys Glu Glu Tyr Glu Ala Ala Arg Thr Glu Leu Gln  
 245 250 255

Glu Glu Ile Ala Arg Gly Ala Ser Ile Gln Asp Ile Arg Ala Arg Leu  
 260 265 270

Thr Lys Thr Asn Asp Lys Ser Gln Ser Lys Glu Glu Pro Leu His Val  
 275 280 285

13.

Thr Lys Ser Glu Ile Pro Asp Asp Leu Ala Gln Ala Gln Ala Tyr Ile  
 290 295 300

Arg Trp Glu Lys Ala Gly Lys Pro Asn Tyr Pro Pro Glu Lys Gln Ile  
 305 310 315 320

Glu Glu Leu Glu Glu Ala Arg Arg Glu Leu Gln Leu Glu Leu Glu Lys  
 325 330 335

Gly Ile Thr Leu Asp Glu Leu Arg Lys Lys Ile Thr Lys Gly Glu Ile  
 340 345 350

Lys Thr Lys Ala Glu Lys His Val Lys Arg Ser Ser Phe Ala Val Glu  
 355 360 365

Arg Ile Gln Arg Lys Lys Arg Asp Phe Gly Gln Leu Ile Asn Lys Tyr  
 370 375 380

Pro Ser Ser Pro Ala Val Gln Val Gln Lys Val Leu Glu Glu Pro Pro  
 385 390 395 400

Ala Leu Ser Lys Ile Lys Leu Tyr Ala Lys Glu Lys Glu Glu Gln Ile  
 405 410 415

Asp Asp Pro Ile Leu Asn Lys Lys Ile Phe Lys Val Asp Asp Gly Glu  
 420 425 430

Leu Leu Val Leu Val Ala Lys Ser Ser Gly Lys Thr Lys Val His Ile  
 435 440 445

Ala Thr Asp Leu Asn Gln Pro Ile Thr Leu His Trp Ala Leu Ser Lys  
 450 455 460

Ser Arg Gly Glu Trp Met Val Pro Pro Ser Ser Ile Leu Pro Pro Gly  
 465 470 475 480

Ser Ile Ile Leu Asp Lys Ala Ala Glu Thr Pro Phe Ser Ala Ser Ser  
 485 490 495

Ser Asp Gly Leu Thr Ser Lys Val Gln Ser Leu Asp Ile Val Ile Glu  
 500 505 510

Asp Gly Asn Phe Val Gly Met Pro Phe Val Leu Leu Ser Gly Glu Lys  
 515 520 525

## 14.

Trp Ile Lys Asn Gln Gly Ser Asp Phe Tyr Val Asp Phe Ser Ala Ala  
 530 535 540

Ser Lys Leu Ala Leu Lys Ala Ala Gly Asp Gly Ser Gly Thr Ala Lys  
 545 550 555 560

Ser Leu Leu Asp Lys Ile Ala Asp Met Glu Ser Glu Ala Gln Lys Ser  
 565 570 575

Phe Met His Arg Phe Asn Ile Ala Ala Asp Leu Ile Glu Asp Ala Thr  
 580 585 590

Ser Ala Gly Glu Leu Gly Phe Thr Gly Ile Leu Val Trp Met Arg Phe  
 595 600 605

Met Ala Thr Arg Gln Leu Ile Trp Asn Lys Asn Tyr Asn Val Lys Pro  
 610 615 620

Arg Glu Ile Ser Lys Ala Gln Asp Arg Leu Thr Asp Leu Leu Gln Asn  
 625 630 635 640

Ala Phe Thr Ser His Pro Gln Tyr Arg Glu Ile Leu Arg Met Ile Met  
 645 650 655

Ser Thr Val Gly Arg Gly Gly Glu Gly Asp Val Gly Gln Arg Ile Arg  
 660 665 670

Asp Glu Ile Leu Val Ile Gln Arg Lys Asn Asp Cys Lys Gly Gly Met  
 675 680 685

Met Glu Glu Trp His Gln Lys Leu His Asn Asn Thr Ser Pro Asp Asp  
 690 695 700

Val Val Ile Cys Gln Ala Leu Ile Asp Tyr Ile Lys Ser Asp Phe Asp  
 705 710 715 720

Leu Gly Val Tyr Trp Lys Thr Leu Asn Glu Asn Gly Ile Thr Lys Glu  
 725 730 735

Arg Leu Leu Ser Tyr Asp Arg Ala Ile His Ser Glu Pro Asn Phe Arg  
 740 745 750

Gly Asp Gln Lys Asn Gly Leu Leu Arg Asp Leu Gly His Tyr Met Arg  
 755 760 765

## 15.

Thr Leu Lys Ala Val His Ser Gly Ala Asp Leu Glu Ser Ala Ile Ala  
 770 775 780

Asn Cys Met Gly Tyr Lys Thr Glu Gly Glu Gly Phe Met Val Gly Val  
 785 790 795 800

Gln Ile Asn Pro Val Ser Gly Leu Pro Ser Gly Phe Gln Gly Leu Leu  
 805 810 815

His Phe Val Leu Asp His Val Glu Asp Lys Asn Val Glu Thr Leu Leu  
 820 825 830

Glu Gly Leu Leu Glu Ala Arg Glu Glu Leu Arg Pro Leu Leu Leu Lys  
 835 840 845

Pro Asn Asn Arg Leu Lys Asp Leu Leu Phe Leu Asp Ile Ala Leu Asp  
 850 855 860

Ser Thr Val Arg Thr Ala Val Glu Arg Gly Tyr Glu Glu Leu Asn Asn  
 865 870 875 880

Ala Asn Pro Glu Lys Ile Met Tyr Phe Ile Ser Leu Val Leu Glu Asn  
 885 890 895

Leu Ala Leu Ser Val Asp Asp Asn Glu Asp Leu Val Tyr Cys Leu Lys  
 900 905 910

Gly Trp Asn Gln Ala Leu Ser Met Ser Asn Gly Gly Asp Asn His Trp  
 915 920 925

Ala Leu Phe Ala Lys Ala Val Leu Asp Arg Ile Arg Leu Ala Leu Ala  
 930 935 940

Ser Lys Ala Glu Trp Tyr His His Leu Leu Gln Pro Ser Ala Glu Tyr  
 945 950 955 960

Leu Gly Ser Ile Leu Gly Val Asp Gln Trp Ala Leu Asn Ile Phe Thr  
 965 970 975

Glu Glu Ile Ile Arg Ala Gly Ser Ala Ala Ser Leu Ser Ser Leu Leu  
 980 985 990

Asn Arg Leu Asp Pro Val Leu Arg Lys Thr Ala Asn Leu Gly Ser Trp  
 995 1000 1005

Gln Ile	Ile Ser Pro Val Glu	Ala Val Gly Tyr Val	Val Val Val
1010	1015	1020	
Asp Glu	Leu Leu Ser Val Gln	Asn Glu Ile Tyr Lys	Lys Pro Thr
1025	1030	1035	
Ile Leu	Val Ala Asn Ser Val	Lys Gly Glu Glu Glu	Ile Pro Asp
1040	1045	1050	
Gly Ala	Val Ala Leu Ile Thr	Pro Asp Met Pro Asp	Val Leu Ser
1055	1060	1065	
His Val	Ser Val Arg Ala Arg	Asn Gly Lys Val Cys	Phe Ala Thr
1070	1075	1080	
Cys Phe	Asp Pro Asn Ile Leu	Ala Asp Leu Gln Ala	Lys Glu Gly
1085	1090	1095	
Arg Ile	Leu Leu Leu Lys Pro	Thr Pro Ser Asp Ile	Ile Tyr Ser
1100	1105	1110	
Glu Val	Asn Glu Ile Glu Leu	Gln Ser Ser Ser Asn	Leu Val Glu
1115	1120	1125	
Ala Glu	Thr Ser Ala Thr Leu	Arg Leu Val Lys Lys	Gln Phe Gly
1130	1135	1140	
Gly Cys	Tyr Ala Ile Ser Ala	Asp Glu Phe Thr Ser	Glu Met Val
1145	1150	1155	
Gly Ala	Lys Ser Arg Asn Ile	Ala Tyr Leu Lys Gly	Lys Val Pro
1160	1165	1170	
Ser Ser	Val Gly Ile Pro Thr	Ser Val Ala Leu Pro	Phe Gly Val
1175	1180	1185	
Phe Glu	Lys Val Leu Ser Asp	Asp Ile Asn Gln Gly	Val Ala Lys
1190	1195	1200	
Glu Leu	Gln Ile Leu Thr Lys	Lys Leu Ser Glu Gly	Asp Phe Ser
1205	1210	1215	
Ala Leu	Gly Glu Ile Arg Thr	Thr Ile Leu Asp Leu	Ser Ala Pro
1220	1225	1230	

Ala Gln	Leu Val Lys Glu Leu	Lys Glu Lys Met Gln	Gly Ser Gly
1235	1240	1245	
Met Pro	Trp Pro Gly Asp Glu	Gly Pro Lys Arg Trp	Glu Gln Ala
1250	1255	1260	
Trp Met	Ala Ile Lys Lys Val	Trp Ala Ser Lys Trp	Asn Glu Arg
1265	1270	1275	
Ala Tyr	Phe Ser Thr Arg Lys	Val Lys Leu Asp His	Asp Tyr Leu
1280	1285	1290	
Cys Met	Ala Val Leu Val Gln	Glu Ile Ile Asn Ala	Asp Tyr Ala
1295	1300	1305	
Phe Val	Ile His Thr Thr Asn	Pro Ser Ser Gly Asp	Asp Ser Glu
1310	1315	1320	
Ile Tyr	Ala Glu Val Val Arg	Gly Leu Gly Glu Thr	Leu Val Gly
1325	1330	1335	
Ala Tyr	Pro Gly Arg Ala Leu	Ser Phe Ile Cys Lys	Lys Lys Asp
1340	1345	1350	
Leu Asn	Ser Pro Gln Val Leu	Gly Tyr Pro Ser Lys	Pro Ile Gly
1355	1360	1365	
Leu Phe	Ile Lys Arg Ser Ile	Ile Phe Arg Ser Asp	Ser Asn Gly
1370	1375	1380	
Glu Asp	Leu Glu Gly Tyr Ala	Gly Ala Gly Leu Tyr	Asp Ser Val
1385	1390	1395	
Pro Met	Asp Glu Glu Glu Lys	Val Val Ile Asp Tyr	Ser Ser Asp
1400	1405	1410	
Pro Leu	Ile Thr Asp Gly Asn	Phe Arg Gln Thr Ile	Leu Ser Asn
1415	1420	1425	
Ile Ala	Arg Ala Gly His Ala	Ile Glu Glu Leu Tyr	Gly Ser Pro
1430	1435	1440	
Gln Asp	Ile Glu Gly Val Val	Arg Asp Gly Lys Ile	Tyr Val Val
1445	1450	1455	



## 18.

Gln Thr Arg Pro Gln Met  
1460

<210> 7  
<211> 1641  
<212> DNA  
<213> Solanum tuberosum

<220>  
<221> CDS  
<222> (1)..(1638)  
<223>

<400> 7  
atg aag cac agt tca gct att tcc gct gtt ttg acc gat gac aat tcg 48  
Met Lys His Ser Ser Ala Ile Ser Ala Val Leu Thr Asp Asp Asn Ser  
1 5 10 15  
aca atg gca ccc cta gag gaa gat gtc aag act gaa aat att ggc ctc 96  
Thr Met Ala Pro Leu Glu Glu Asp Val Lys Thr Glu Asn Ile Gly Leu  
20 25 30  
cta aat ttg gat cca act ttg gaa cct tat cta gat cac ttc aga cac 144  
Leu Asn Leu Asp Pro Thr Leu Glu Pro Tyr Leu Asp His Phe Arg His  
35 40 45  
aga atg aag aga tat gtg gat cag aaa atg ctc att gaa aaa tat gag 192  
Arg Met Lys Arg Tyr Val Asp Gln Lys Met Leu Ile Glu Lys Tyr Glu  
50 55 60  
gga ccc ctt gag gaa ttt gct caa ggt tat tta aaa ttt gga ttc aac 240  
Gly Pro Leu Glu Glu Phe Ala Gln Gly Tyr Leu Lys Phe Gly Phe Asn  
65 70 75 80  
agg gaa gat ggt tgc ata gtc tat cgt gaa tgg gct cct gct gct cag 288  
Arg Glu Asp Gly Cys Ile Val Tyr Arg Glu Trp Ala Pro Ala Ala Gln  
85 90 95  
gaa gca gaa gtt att ggc gat ttc aat gga tgg aac ggt tct aac cac 336  
Glu Ala Glu Val Ile Gly Asp Phe Asn Gly Trp Asn Gly Ser Asn His  
100 105 110  
atg atg gag aag gac cag ttt ggt gtt tgg agt att aga att cct gat 384  
Met Met Glu Lys Asp Gln Phe Gly Val Trp Ser Ile Arg Ile Pro Asp  
115 120 125  
gtt gac agt aag cca gtc att cca cac aac tcc aga gtt aag ttt cgt 432  
Val Asp Ser Lys Pro Val Ile Pro His Asn Ser Arg Val Lys Phe Arg  
130 135 140  
ttc aaa cat ggt aat gga gtg tgg gta gat cgt atc cct gct tgg ata 480  
Phe Lys His Gly Asn Val Trp Val Asp Arg Ile Pro Ala Trp Ile  
145 150 155 160  
aag tat gcc act gca gac gcc aca aag ttt gca gca cca tat gat ggt 528  
Lys Tyr Ala Thr Ala Asp Ala Thr Lys Phe Ala Ala Pro Tyr Asp Gly  
165 170 175

## 19.

gtc tac tgg gac cca cca cct tca gaa agg tac cac ttc aaa tac cct	576
Val Tyr Trp Asp Pro Pro Pro Ser Glu Arg Tyr His Phe Lys Tyr Pro	
180 185 190	
cgc cct ccc aaa ccc cga gcc cca cga atc tac gaa gca cat gtc ggc	624
Arg Pro Pro Lys Pro Arg Ala Pro Arg Ile Tyr Glu Ala His Val Gly	
195 200 205	
atg agc agc tct gag cca cgt gta aat tgc tat cgt gag ttt gca gat	672
Met Ser Ser Ser Glu Pro Arg Val Asn Ser Tyr Arg Glu Phe Ala Asp	
210 215 220	
gat gtt tta cct cgg att aag gca aat aac tat aat act gtc cag ttg	720
Asp Val Leu Pro Arg Ile Lys Ala Asn Asn Tyr Asn Thr Val Gln Leu	
225 230 235 240	
atg gcc ata atg gaa cat tct tac tat gga tca ttt gga tat cat gtt	768
Met Ala Ile Met Glu His Ser Tyr Tyr Gly Ser Phe Gly Tyr His Val	
245 250 255	
aca aac ttt ttt gct gtg agc aat aga tat gga aac ccg gag gac cta	816
Thr Asn Phe Phe Ala Val Ser Asn Arg Tyr Gly Asn Pro Glu Asp Leu	
260 265 270	
aag tat ctg ata gat aaa gca cat agc ttg ggt tta cag gtt ctg gtg	864
Lys Tyr Leu Ile Asp Lys Ala His Ser Leu Gly Leu Gln Val Leu Val	
275 280 285	
gat gta gtt cac agt cat gca agc aat aat gtc act gat ggc ctc aat	912
Asp Val Val His Ser His Ala Ser Asn Asn Val Thr Asp Gly Leu Asn	
290 295 300	
ggc ttt gat att ggc caa ggt tct caa gaa tcc tac ttt cat gct gga	960
Gly Phe Asp Ile Gly Gln Gly Ser Gln Glu Ser Tyr Phe His Ala Gly	
305 310 315 320	
gag cga ggg tac cat aag ttg tgg gat agc agg ctg ttc aac tat gcc	1008
Glu Arg Gly Tyr His Lys Leu Trp Asp Ser Arg Leu Phe Asn Tyr Ala	
325 330 335	
aat tgg gag gtt ctt cgt ttc ctt ctt tcc aac ttg agg tgg tgg cta	1056
Asn Trp Glu Val Leu Arg Phe Leu Leu Ser Asn Leu Arg Trp Trp Leu	
340 345 350	
gaa gag tat aac ttt gac gga ttt cga ttt gat gga ata act tct atg	1104
Glu Glu Tyr Asn Phe Asp Gly Phe Arg Phe Asp Gly Ile Thr Ser Met	
355 360 365	
ctg tat gtt cat cat gga atc aat atg gga ttt aca gga aac tat aat	1152
Leu Tyr Val His His Gly Ile Asn Met Gly Phe Thr Gly Asn Tyr Asn	
370 375 380	
gag tat ttc agc gag gct aca gat gtt gat gct gtg gtc tat tta atg	1200
Glu Tyr Phe Ser Glu Ala Thr Asp Val Asp Ala Val Val Tyr Leu Met	
385 390 395 400	
ttg gcc aat aat ctg att cac aag att ttc cca gac gca act gtt att	1248
Leu Ala Asn Asn Leu Ile His Lys Ile Phe Pro Asp Ala Thr Val Ile	
405 410 415	

## 20.

gcc gaa gat gtt tct ggt atg ccg ggc ctt agc cgg cct gtt tct gag 1296  
 Ala Glu Asp Val Ser Gly Met Pro Gly Leu Ser Arg Pro Val Ser Glu  
 420 425 430

gga gga att ggt ttt gat tac cgc ctg gca atg gca atc cca gat aag 1344  
 Gly Gly Ile Gly Phe Asp Tyr Arg Leu Ala Met Ala Ile Pro Asp Lys  
 435 440 445

tgg ata gat tat tta aag aat aag aat gat gaa gat tgg tcc atg aag 1392  
 Trp Ile Asp Tyr Leu Lys Asn Lys Asn Asp Glu Asp Trp Ser Met Lys  
 450 455 460

gaa gta aca tcg agt ttg aca aat agg aga tat aca gag aag tgt ata 1440  
 Glu Val Thr Ser Ser Leu Thr Asn Arg Arg Tyr Thr Glu Lys Cys Ile  
 465 470 475 480

gca tat gcg gag agc cat gat cag tct att gtc ggt gac aag acc att 1488  
 Ala Tyr Ala Glu Ser His Asp Gln Ser Ile Val Gly Asp Lys Thr Ile  
 485 490 495

gca ttt ctc cta atg gac aaa gag atg tat tct ggc atg tct tgc ttg 1536  
 Ala Phe Leu Leu Met Asp Lys Glu Met Tyr Ser Gly Met Ser Cys Leu  
 500 505 510

aca gat gct tct cct gtt gtt gat cga gga att gcg ctt cac aag atg 1584  
 Thr Asp Ala Ser Pro Val Val Asp Arg Gly Ile Ala Leu His Lys Met  
 515 520 525

atc cat ttt ttt cac aat ggc ctt ggg agg aga ggg gta cct caa ttt 1632  
 Ile His Phe Phe His Asn Gly Leu Gly Arg Arg Gly Val Pro Gln Phe  
 530 535 540

cat ggg taa 1641  
 His Gly  
 545

<210> 8  
 <211> 546  
 <212> PRT  
 <213> Solanum tuberosum

<400> 8

Met Lys His Ser Ser Ala Ile Ser Ala Val Leu Thr Asp Asp Asn Ser  
 1 5 10 15

Thr Met Ala Pro Leu Glu Glu Asp Val Lys Thr Glu Asn Ile Gly Leu  
 20 25 30

Leu Asn Leu Asp Pro Thr Leu Glu Pro Tyr Leu Asp His Phe Arg His  
 35 40 45

Arg Met Lys Arg Tyr Val Asp Gln Lys Met Leu Ile Glu Lys Tyr Glu  
 50 55 60

## 21

Gly Pro Leu Glu Glu Phe Ala Gln Gly Tyr Leu Lys Phe Gly Phe Asn  
 65 70 75 80

Arg Glu Asp Gly Cys Ile Val Tyr Arg Glu Trp Ala Pro Ala Ala Gln  
 85 90 95

Glu Ala Glu Val Ile Gly Asp Phe Asn Gly Trp Asn Gly Ser Asn His  
 100 105 110

Met Met Glu Lys Asp Gln Phe Gly Val Trp Ser Ile Arg Ile Pro Asp  
 115 120 125

Val Asp Ser Lys Pro Val Ile Pro His Asn Ser Arg Val Lys Phe Arg  
 130 135 140

Phe Lys His Gly Asn Gly Val Trp Val Asp Arg Ile Pro Ala Trp Ile  
 145 150 155 160

Lys Tyr Ala Thr Ala Asp Ala Thr Lys Phe Ala Ala Pro Tyr Asp Gly  
 165 170 175

Val Tyr Trp Asp Pro Pro Pro Ser Glu Arg Tyr His Phe Lys Tyr Pro  
 180 185 190

Arg Pro Pro Lys Pro Arg Ala Pro Arg Ile Tyr Glu Ala His Val Gly  
 195 200 205

Met Ser Ser Ser Glu Pro Arg Val Asn Ser Tyr Arg Glu Phe Ala Asp  
 210 215 220

Asp Val Leu Pro Arg Ile Lys Ala Asn Asn Tyr Asn Thr Val Gln Leu  
 225 230 235 240

Met Ala Ile Met Glu His Ser Tyr Tyr Gly Ser Phe Gly Tyr His Val  
 245 250 255

Thr Asn Phe Phe Ala Val Ser Asn Arg Tyr Gly Asn Pro Glu Asp Leu  
 260 265 270

Lys Tyr Leu Ile Asp Lys Ala His Ser Leu Gly Leu Gln Val Leu Val  
 275 280 285

Asp Val Val His Ser His Ala Ser Asn Asn Val Thr Asp Gly Leu Asn  
 290 295 300

Gly Phe Asp Ile Gly Gln Gly Ser Gln Glu Ser Tyr Phe His Ala Gly  
 305 310 315 320

Glu Arg Gly Tyr His Lys Leu Trp Asp Ser Arg Leu Phe Asn Tyr Ala  
 325 330 335

Asn Trp Glu Val Leu Arg Phe Leu Leu Ser Asn Leu Arg Trp Trp Leu  
 340 345 350

Glu Glu Tyr Asn Phe Asp Gly Phe Arg Phe Asp Gly Ile Thr Ser Met  
 355 360 365

Leu Tyr Val His His Gly Ile Asn Met Gly Phe Thr Gly Asn Tyr Asn  
 370 375 380

Glu Tyr Phe Ser Glu Ala Thr Asp Val Asp Ala Val Val Tyr Leu Met  
 385 390 395 400

Leu Ala Asn Asn Leu Ile His Lys Ile Phe Pro Asp Ala Thr Val Ile  
 405 410 415

Ala Glu Asp Val Ser Gly Met Pro Gly Leu Ser Arg Pro Val Ser Glu  
 420 425 430

Gly Gly Ile Gly Phe Asp Tyr Arg Leu Ala Met Ala Ile Pro Asp Lys  
 435 440 445

Trp Ile Asp Tyr Leu Lys Asn Lys Asn Asp Glu Asp Trp Ser Met Lys  
 450 455 460

Glu Val Thr Ser Ser Leu Thr Asn Arg Arg Tyr Thr Glu Lys Cys Ile  
 465 470 475 480

Ala Tyr Ala Glu Ser His Asp Gln Ser Ile Val Gly Asp Lys Thr Ile  
 485 490 495

Ala Phe Leu Leu Met Asp Lys Glu Met Tyr Ser Gly Met Ser Cys Leu  
 500 505 510

Thr Asp Ala Ser Pro Val Val Asp Arg Gly Ile Ala Leu His Lys Met  
 515 520 525

Ile His Phe Phe His Asn Gly Leu Gly Arg Arg Gly Val Pro Gln Phe  
 530 535 540

His Gly  
545

<210> 9  
<211> 822  
<212> PRT  
<213> Zea mays

<400> 9

Leu Cys Leu Val Ser Pro Ser Ser Ser Pro Thr Pro Leu Pro Pro Pro  
1 5 10 15

Arg Arg Ser Arg Ser His Ala Asp Arg Ala Ala Pro Pro Gly Ile Ala  
20 25 30

Gly Gly Gly Asn Val Arg Leu Ser Val Leu Ser Val Gln Cys Lys Ala  
35 40 45

Arg Arg Ser Gly Val Arg Lys Val Lys Ser Lys Phe Ala Thr Ala Ala  
50 55 60

Thr Val Gln Glu Asp Lys Thr Met Ala Thr Ala Lys Gly Asp Val Asp  
65 70 75 80

His Leu Pro Ile Tyr Asp Leu Asp Pro Lys Leu Glu Ile Phe Lys Asp  
85 90 95

His Phe Arg Tyr Arg Met Lys Arg Phe Leu Glu Gln Lys Gly Ser Ile  
100 105 110

Glu Glu Asn Glu Gly Ser Leu Glu Ser Phe Ser Lys Gly Tyr Leu Lys  
115 120 125

Phe Gly Ile Asn Thr Asn Glu Asp Gly Thr Val Tyr Arg Glu Trp Ala  
130 135 140

Pro Ala Ala Gln Glu Ala Glu Leu Ile Gly Asp Phe Asn Asp Trp Asn  
145 150 155 160

Gly Ala Asn His Lys Met Glu Lys Asp Lys Phe Gly Val Trp Ser Ile  
165 170 175

Lys Ile Asp His Val Lys Gly Lys Pro Ala Ile Pro His Asn Ser Lys  
180 185 190

## 24

Val Lys Phe Arg Phe Leu His Gly Gly Val Trp Val Asp Arg Ile Pro  
 195 200 205

Ala Leu Ile Arg Tyr Ala Thr Val Asp Ala Ser Lys Phe Gly Ala Pro  
 210 215 220

Tyr Asp Gly Val His Trp Asp Pro Pro Ala Ser Glu Arg Tyr Thr Phe  
 225 230 235 240

Lys His Pro Arg Pro Ser Lys Pro Ala Ala Pro Arg Ile Tyr Glu Ala  
 245 250 255

His Val Gly Met Ser Gly Glu Lys Pro Ala Val Ser Thr Tyr Arg Glu  
 260 265 270

Phe Ala Asp Asn Val Leu Pro Arg Ile Arg Ala Asn Asn Tyr Asn Thr  
 275 280 285

Val Gln Leu Met Ala Val Met Glu His Ser Tyr Tyr Ala Ser Phe Gly  
 290 295 300

Tyr His Val Thr Asn Phe Phe Ala Val Ser Ser Arg Ser Gly Thr Pro  
 305 310 315 320

Glu Asp Leu Lys Tyr Leu Val Asp Lys Ala His Ser Leu Gly Leu Arg  
 325 330 335

Val Leu Met Asp Val Val His Ser His Ala Ser Asn Asn Val Thr Asp  
 340 345 350

Gly Leu Asn Gly Tyr Asp Val Gly Gln Ser Thr Gln Glu Ser Tyr Phe  
 355 360 365

His Ala Gly Asp Arg Gly Tyr His Lys Leu Trp Asp Ser Arg Leu Phe  
 370 375 380

Asn Tyr Ala Asn Trp Glu Val Leu Arg Phe Leu Leu Ser Asn Leu Arg  
 385 390 395 400

Tyr Trp Leu Asp Glu Phe Met Phe Asp Gly Phe Arg Phe Asp Gly Val  
 405 410 415

Thr Ser Met Leu Tyr His His His Gly Ile Asn Val Gly Phe Thr Gly  
 420 425 430

## 25

Asn Tyr Gln Glu Tyr Phe Ser Leu Asp Thr Ala Val Asp Ala Val Val  
 435 440 445

Tyr Met Met Leu Ala Asn His Leu Met His Lys Leu Leu Pro Glu Ala  
 450 455 460

Thr Val Val Ala Glu Asp Val Ser Gly Met Pro Val Leu Cys Arg Pro  
 465 470 475 480

Val Asp Glu Gly Gly Val Gly Phe Asp Tyr Arg Leu Ala Met Ala Ile  
 485 490 495

Pro Asp Arg Trp Ile Asp Tyr Leu Lys Asn Lys Asp Asp Ser Glu Trp  
 500 505 510

Ser Met Gly Glu Ile Ala His Thr Leu Thr Asn Arg Arg Tyr Thr Glu  
 515 520 525

Lys Cys Ile Ala Tyr Ala Glu Ser His Asp Gln Ser Ile Val Gly Asp  
 530 535 540

Lys Thr Ile Ala Phe Leu Leu Met Asp Lys Glu Met Tyr Thr Gly Met  
 545 550 555 560

Ser Asp Leu Gln Pro Ala Ser Pro Thr Ile Asp Arg Gly Ile Ala Leu  
 565 570 575

Gln Lys Met Ile His Phe Ile Thr Met Ala Leu Gly Gly Asp Gly Tyr  
 580 585 590

Leu Asn Phe Met Gly Asn Glu Phe Gly His Pro Glu Trp Ile Asp Phe  
 595 600 605

Pro Arg Glu Gly Asn Asn Trp Ser Tyr Asp Lys Cys Arg Arg Gln Trp  
 610 615 620

Ser Leu Val Asp Thr Asp His Leu Arg Tyr Lys Tyr Met Asn Ala Phe  
 625 630 635 640

Asp Gln Ala Met Asn Ala Leu Asp Glu Arg Phe Ser Phe Leu Ser Ser  
 645 650 655

Ser Lys Gln Ile Val Ser Asp Met Asn Asp Glu Glu Lys Val Ile Val  
 660 665 670



## 26.

Phe Glu Arg Gly Asp Leu Val Phe Val Phe Asn Phe His Pro Lys Lys  
           675                          680                          685

Thr Tyr Glu Gly Tyr Lys Val Gly Cys Asp Leu Pro Gly Lys Tyr Arg  
       690                          695                          700

Val Ala Leu Asp Ser Asp Ala Leu Val Phe Gly Gly His Gly Arg Val  
       705                          710                          715                          720

Gly His Asp Val Asp His Phe Thr Ser Pro Glu Gly Val Pro Gly Val  
                           725                          730                          735

Pro Glu Thr Asn Phe Asn Asn Arg Pro Asn Ser Phe Lys Val Leu Ser  
                           740                          745                          750

Pro Pro Arg Thr Cys Val Ala Tyr Tyr Arg Val Asp Glu Ala Gly Ala  
                           755                          760                          765

Gly Arg Arg Leu His Ala Lys Ala Glu Thr Gly Lys Thr Ser Pro Ala  
       770                          775                          780

Glu Ser Ile Asp Val Lys Ala Ser Arg Ala Ser Ser Lys Glu Asp Lys  
       785                          790                          795                          800

Glu Ala Thr Ala Gly Gly Lys Lys Gly Trp Lys Phe Ala Arg Gln Pro  
                           805                          810                          815

Ser Asp Gln Asp Thr Lys  
                           820

<210> 10  
 <211> 814  
 <212> PRT  
 <213> Zea mays

<400> 10

Asp Leu Pro Ser Val Leu Phe Arg Arg Lys Asp Ala Phe Ser Arg Thr  
   1                          5                          10                          15

Val Leu Ser Cys Ala Gly Ala Pro Gly Lys Val Leu Val Pro Gly Gly  
           20                          25                          30

Gly Ser Asp Asp Leu Leu Ser Ser Ala Glu Pro Val Val Asp Thr Gln  
       35                          40                          45

Pro Glu Glu Leu Gln Ile Pro Glu Ala Glu Leu Thr Val Glu Lys Thr

27

50		55		60											
Ser 65	Ser 66	Ser 67	Pro 68	Thr 69	Gln 70	Thr 71	Thr 72	Ser 73	Ala 74	Val 75	Ala 76	Glu 77	Ala 78	Ser 79	Ser 80
Gly 81	Val 82	Glu 83	Ala 84	Glu 85	Glu 86	Arg 87	Pro 88	Glu 89	Leu 90	Ser 91	Glu 92	Val 93	Ile 94	Gly 95	Val 96
Gly 97	Gly 98	Thr 99	Gly 100	Gly 101	Thr 102	Lys 103	Ile 104	Asp 105	Gly 106	Ala 107	Gly 108	Ile 109	Lys 110	Ala 111	Lys 112
Ala 113	Pro 114	Leu 115	Val 116	Glu 117	Glu 118	Lys 119	Pro 120	Arg 121	Val 122	Ile 123	Pro 124	Pro 125	Pro 126	Gly 127	Asp 128
Gly 129	Gln 130	Arg 131	Ile 132	Tyr 133	Glu 134	Ile 135	Asp 136	Pro 137	Met 138	Leu 139	Glu 140	Gly 141	Phe 142	Arg 143	Gly 144
His 145	Leu 146	Asp 147	Tyr 148	Arg 149	Tyr 150	Ser 151	Glu 152	Tyr 153	Lys 154	Arg 155	Leu 156	Arg 157	Ala 158	Ala 159	Ile 160
Asp 161	Gln 162	His 163	Glu 164	Gly 165	Gly 166	Leu 167	Asp 168	Ala 169	Phe 170	Ser 171	Arg 172	Gly 173	Tyr 174	Glu 175	Lys 176
Leu 177	Gly 178	Phe 179	Thr 180	Arg 181	Ser 182	Ala 183	Glu 184	Gly 185	Ile 186	Thr 187	Tyr 188	Arg 189	Glu 190	Trp 191	Ala 192
Pro 193	Gly 194	Ala 195	Tyr 196	Ser 197	Ala 198	Ala 199	Leu 200	Val 201	Gly 202	Asp 203	Phe 204	Asn 205	Asn 206	Trp 207	Asn 208
Pro 209	Asn 210	Ala 211	Asp 212	Ala 213	Met 214	Ala 215	Arg 216	Asn 217	Glu 218	Tyr 219	Gly 220	Val 221	Trp 222	Glu 223	Ile 224
Phe 225	Leu 226	Pro 227	Asn 228	Asn 229	Ala 230	Asp 231	Gly 232	Ser 233	Pro 234	Ala 235	Ile 236	Pro 237	His 238	Gly 239	Ser 240
Arg 241	Val 242	Lys 243	Ile 244	Arg 245	Met 246	Asp 247	Thr 248	Pro 249	Ser 250	Gly 251	Val 252	Lys 253	Asp 254	Ser 255	Ile 256
Pro 257	Ala 258	Trp 259	Ile 260	Lys 261	Phe 262	Ser 263	Val 264	Gln 265	Ala 266	Pro 267	Gly 268	Glu 269	Ile 270	Pro 271	Tyr 272
Asn 273	Gly 274	Ile 275	Tyr 276	Tyr 277	Asp 278	Pro 279	Pro 280	Glu 281	Glu 282	Glu 283	Lys 284	Tyr 285	Val 286	Phe 287	Lys 288
His 289	Pro 290	Gln 291	Pro 292	Lys 293	Arg 294	Pro 295	Lys 296	Ser 297	Leu 298	Arg 299	Ile 300	Tyr 301	Glu 302	Ser 303	His 304

28.

290	295	300
Val Gly Met Ser Ser Pro Glu Pro Lys Ile Asn Thr Tyr Ala Asn Phe 305 310 315 320		
Arg Asp Glu Val Leu Pro Arg Ile Lys Lys Leu Gly Tyr Asn Ala Val 325 330 335		
Gln Ile Met Ala Ile Gln Glu His Ser Tyr Tyr Ala Ser Phe Gly Tyr 340 345 350		
His Val Thr Asn Phe Phe Ala Pro Ser Ser Arg Phe Gly Thr Pro Glu 355 360 365		
Asp Leu Lys Ser Leu Ile Asp Lys Ala His Glu Leu Gly Leu Leu Val 370 375 380		
Leu Met Asp Ile Val His Ser His Ser Ser Asn Asn Thr Leu Asp Gly 385 390 395 400		
Leu Asn Gly Phe Asp Gly Thr Asp Thr His Tyr Phe His Gly Gly Pro 405 410 415		
Arg Gly His His Trp Met Trp Asp Ser Arg Leu Phe Asn Tyr Gly Ser 420 425 430		
Trp Glu Val Leu Arg Phe Leu Leu Ser Asn Ala Arg Trp Trp Leu Glu 435 440 445		
Glu Tyr Lys Phe Asp Gly Phe Arg Phe Asp Gly Val Thr Ser Met Met 450 455 460		
Tyr Thr His His Gly Leu Gln Val Thr Phe Thr Gly Asn Tyr Gly Glu 465 470 475 480		
Tyr Phe Gly Phe Ala Thr Asp Val Asp Ala Val Val Tyr Leu Met Leu 485 490 495		
Val Asn Asp Leu Ile Arg Gly Leu Tyr Pro Glu Ala Val Ser Ile Gly 500 505 510		
Glu Asp Val Ser Gly Met Pro Thr Phe Cys Ile Pro Val Gln Asp Gly 515 520 525		
Gly Val Gly Phe Asp Tyr Arg Leu His Met Ala Val Pro Asp Lys Trp		

## 29.

530	535	540
Ile Glu Leu Leu Lys Gln Ser Asp Glu Tyr Trp Glu Met Gly Asp Ile		
545	550	555 560
Val His Thr Leu Thr Asn Arg Arg Trp Leu Glu Lys Cys Val Thr Tyr		
	565	570 575
Cys Glu Ser His Asp Gln Ala Leu Val Gly Asp Lys Thr Ile Ala Phe		
	580	585 590
Trp Leu Met Asp Lys Asp Met Tyr Asp Phe Met Ala Leu Asp Arg Pro		
	595	600 605
Ser Thr Pro Arg Ile Asp Arg Gly Ile Ala Leu His Lys Met Ile Arg		
	610	615 620
Leu Val Thr Met Gly Leu Gly Gly Glu Gly Tyr Leu Asn Phe Met Gly		
	625	630 635 640
Asn Glu Phe Gly His Pro Glu Trp Ile Asp Phe Pro Arg Gly Pro Gln		
	645	650 655
Ser Leu Pro Asn Gly Ser Val Ile Pro Gly Asn Asn Asn Ser Phe Asp		
	660	665 670
Lys Cys Arg Arg Arg Phe Asp Leu Gly Asp Ala Asp Tyr Leu Arg Tyr		
	675	680 685
Arg Gly Met Gln Glu Phe Asp Gln Ala Met Gln His Leu Glu Gly Lys		
	690	695 700
Tyr Glu Phe Met Thr Ser Asp His Ser Tyr Val Ser Arg Lys His Glu		
	705	710 715 720
Glu Asp Lys Val Ile Ile Phe Glu Arg Gly Asp Leu Val Phe Val Phe		
	725	730 735
Asn Phe His Trp Ser Asn Ser Tyr Phe Asp Tyr Arg Val Gly Cys Phe		
	740	745 750
Lys Pro Gly Lys Tyr Lys Ile Val Leu Asp Ser Asp Asp Gly Leu Phe		
	755	760 765
Gly Gly Phe Ser Arg Leu Asp His Asp Ala Glu Tyr Phe Thr Ala Asp		

30.

770

775

780

Trp Pro His Asp Asn Arg Pro Cys Ser Phe Ser Val Tyr Ala Pro Ser  
785 790 795 800

Arg Thr Ala Val Val Tyr Ala Pro Ala Gly Ala Glu Asp Glu  
805 810

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
19 December 2002 (19.12.2002)

PCT

(10) International Publication Number  
**WO 2002/101059 A3**

(51) International Patent Classification<sup>7</sup>: **C12N 15/82**,  
C07K 14/415, A01H 5/00

(21) International Application Number:  
PCT/EP2002/006265

(22) International Filing Date: 7 June 2002 (07.06.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
101 28 363.6 12 June 2001 (12.06.2001) DE

(71) Applicant: BAYER CROPSCIENCE GMBH [DE/DE];  
Brüningstrasse 50, 65929 Frankfurt (DE).

(72) Inventors: UWER, Ursula; Hasenheide 49, 10967 Berlin  
(DE). FROHBERG, Claus; Apoldaer Strasse 14, 12249  
Berlin (DE). PILLING, Jens; Melchiorstrasse 36, 50670  
Köln (DE). LANDSCHÜTZE, Volker; Gardeschützen-  
weg 51, 12203 Berlin (DE).

(74) Agent: VOSSIUS & PARTNER; Siebertstrasse 4, 81675  
Munich (DE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN,  
YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,  
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent  
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,  
NE, SN, TD, TG).

**Published:**

— with international search report

(88) Date of publication of the international search report:  
23 September 2004

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: TRANSGENIC PLANTS SYNTHESISING HIGH AMYLOSE STARCH

(57) Abstract: Transgenic plant cells and plants are described which synthesise a starch which is modified in comparison with corresponding wild type plant cells and plants. The plant cells and plants described show a reduced activity of R1, BEI and BEII proteins. Furthermore, modified starches as well as methods for their production are described.

WO 2002/101059 A3

# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/EP 02/06265

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C12N15/82 C07K14/415 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C12N C07K A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, SEQUENCE SEARCH, BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	LORBERTH ET AL: "Inhibition of a starch-granule-bound protein leads to modified starch and repression of cold sweetening" NATURE BIOTECHNOLOGY, NATURE PUBLISHING, US, vol. 16, May 1998 (1998-05), pages 473-477, XP002111459 ISSN: 1087-0156 cited in the application	18-22
Y	page 476, right-hand column the whole document  --- -/--	1-17, 23-29

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"Z" document member of the same patent family

Date of the actual completion of the international search

7 May 2003

Date of mailing of the international search report

22/05/2003

Name and mailing address of the ISA

European Patent Office, P.O. Box 5018 Patentamt 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Bucka, A

# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/EP 02/06265

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SCHWALL GERHARD P ET AL: "Production of very-high-amylose potato starch by inhibition of SBE A and B." NATURE BIOTECHNOLOGY, vol. 18, no. 5, May 2000 (2000-05), pages 551-554, XP002239695 ISSN: 1087-0156 cited in the application	18-22
Y	page 554, left-hand column the whole document	1-17, 23-29
X	--- WO 97 11188 A (KOSSMANN JENS ; LORBERTH RUTH (DE); PLANTTEC BIOTECHNOLOGIE GMBH (D) 27 March 1997 (1997-03-27) cited in the application	18-22
Y	claims 1,2,12 the whole document	1-17, 23-29
X	--- DE 196 53 176 A (PLANTTEC BIOTECHNOLOGIE GMBH F) 25 June 1998 (1998-06-25) claims 1,2,12	18-22
Y	the whole document	1-17, 23-29
A	--- AMERICAN SOCIETY OF AGRONOMY-OLSON & FREY EDS: "Chapter 7, Par. 4.1.2.1: Starch and Chapter 7, Par 5.2.4.2: High Amylose Corn (GLOVER & MERTZ)" NUTRITIONAL QUALITY OF CEREAL GRAINS: GENETIC AND AGRONOMIC IMPROVEMENT, XX, XX, 1987, pages 183,201-202,269-271, XP002099180 page 270	1-29
A	--- SHI YON-CHENG ET AL: "Molecular structure of a low-amylopectin starch and other high-amylose maize starches." JOURNAL OF CEREAL SCIENCE, vol. 27, no. 3, May 1998 (1998-05), pages 289-299, XP002239696 ISSN: 0733-5210 cited in the application	1-29
	the whole document -----	



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/06265

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9711188	A	27-03-1997	DE 19534759 A1	20-03-1997
			DE 19547733 A1	26-06-1997
			AU 715944 B2	10-02-2000
			AU 7131396 A	09-04-1997
			CA 2231774 A1	27-03-1997
			WO 9711188 A1	27-03-1997
			EP 0851934 A1	08-07-1998
			HU 9900510 A2	28-05-1999
			JP 11512286 T	26-10-1999
			US 6207880 B1	27-03-2001
			US 2001007155 A1	05-07-2001
DE 19653176	A	25-06-1998	DE 19653176 A1	25-06-1998
			AU 740492 B2	08-11-2001
			AU 5857798 A	15-07-1998
			WO 9827212 A1	25-06-1998
			EP 0950107 A1	20-10-1999
			JP 2001522223 T	13-11-2001